

Received: 12.02.2023
Accepted: 30.06.2023

A – Study Design
B – Data Collection
C – Statistical Analysis
D – Data Interpretation
E – Manuscript Preparation
F – Literature Search
G – Funds Collection

PSYCHOGENIC FACTORS AS A CAUSE OF DEPRESSIVE DISORDERS IN THE ELDERLY

Madina Kassymzhanova^[A,B,C,D,E,F,G]

ORCID: 0000-0001-7102-4171

Natalya Raspopova^[A,B,C,D,E,F]

ORCID: 0000-0001-8456-1822

Department of Psychiatry and Narcology Kazakh National Medical University named after S.D. Asfendiyarov, Republic of Kazakhstan

Background:

Material/ Methods:

Results:

Conclusions:

SUMMARY

Individuals with particular qualities or characteristics are pre-disposed to develop cognitive impairment non dementia. The main aim to study socio-demographic factors and personality traits, that contribute to the development of depressive disorders in elderly patients with cognitive deficits. The purpose of the article was to show that it is important to investigate the socio-psychological factors that can worsen the course of cognitive impairment, which already comes with age.

During the research an observational study involving 111 patients from Kazakhstan with cognitive deficits was conducted: 59 patients, whose depressive disorders had been identified by clinical and psychopathological research using the Hamilton Depression Rating Scale (HRSD) and 52 patients for whom psychogenic depressive disorders had not been defined. The recruitment of the study participants was carried out in the state nursing home for the elderly and disabled. The study included male and female individuals aged 60-74 years with different national and ethnic backgrounds, with mild to moderate cognitive impairment, who wished to participate in the study and signed an informed consent to participate in the study.

This study uncovered the primary factors that increase the chances of individuals with cognitive impairments developing a depressive disorder. These factors include personal attributes of the patients, such as their sex, whether they are married or not, and the quality of their relationship with their family members. In addition, personality traits from before their cognitive decline, such as a tendency for isolation, insecurity, and passivity, were also significant.

This study could have an exceptional value for medicine and psychological workers on the whole and in Republic of Kazakhstan in particular to prevent depressive diseases were mentioned.

Keywords: depressive disorders, cognitive deficits, elderly people, psychological diseases, Kazakhstan

INTRODUCTION

The elderly population aged 60 years or over is steadily growing worldwide. From 2015 to 2050 the proportion of people over 60 in the world's population is expected to increase from 12% to 22%. In absolute terms, this percentage increase translates as an increase from 900 million to 2 billion elderly people. The most common neuropsychiatric disorders in this age group are dementia and depression, affecting ~5% and 7% of the world's older population, respectively, while anxiety disorders account for 3.8% of older people. In addition, ~25% of deaths from self-harm occur in people aged 60 years or older (World Population Ageing, 2017).

Old age is a sensitive period during a person's life and is second only to adolescence in terms of vulnerability (Shah et al., 2016). From the age of 60, the influence of psychogenic factors increases, ultimately triggering the development of mental disorders. Great contribution to the study of depressive disorders in the elderly of Kazakhstan was made by scientists R. G. Ilesheva (1981), G. M. Kudyarova (1989), Zaltsman I. G. (1998) Izmailova N. T. et al. (2000). In their works is noted, that at this age depression is one of the most frequent types of mental pathology. Compared with depression in younger people, the proportion of depressive states with psychomotor retardation decreases. On the contrary, anxious agitation, sometimes reaching raptus, is much more often observed. Mental nerve block is less common but quite often emotional lability, irritability, and gloominess. The theme of depressive experiences is dominated by the ideas of damage and impoverishment and hypochondria. These factors can also aggravate the course and outcome of somatic diseases. Several studies highlight the importance of regular assessment of neuropsychiatric symptoms in patients with cognitive deficits in primary geriatric care services, in particular depression (Kim et al., 2020; Ikezaki et al., 2020; Ginsberg et al., 2019; Romero-Sevilla et al., 2018; Lacour et al., 2017). Indeed, by focusing on identifying depressive disorders in people at risk of developing dementia, medicine workers can help slow the rate of progression of cognitive deficits (Lara, 2017; Vicini Chilovi et al., 2009; Yang et al., 2020; Fischer et al., 2019; Yatawara et al., 2018; Petkus et al., 2017). In addition, there is accumulating evidence of an unfavorable association between anxiety and cognitive decline (Mora, 2013; Jia et al., 2020; Hu et al., 2021b; Zufferey et al., 2020; Watanabe et al., 2020; Rainero et al., 2020). This correlation is bi-directional; anxiety can increase with the progression of neurodegeneration and be an independent factor in the deterioration of cognitive status (Jenkins et al., 2021; Gigi and Papirovitza, 2021; Thancharoen et al., 2020; Seifan et al., 2019; Nolte et al., 2019).

In addition to studying risk factors for the occurrence of cognitive deficits, researchers have also investigated protective factors (Pais et al., 2021; Delfino et al., 2018; Sutin et al., 2018; Johansson et al., 2014). There are many studies on the influence of protective factors on the development of various types of dementia, including having an active social life, participating in physical activity,

maintaining emotional wellness, and controlling the negative effects of stress (Wiegelmann et al., 2021; Sutin et al., 2021b; McGrattan et al., 2021; Krivanek et al., 2021; Fu et al., 2021; Floud et al., 2021). Recently published results from the Singapore Geriatric Study on the Status of Elderly show a correlation between happiness and higher cognitive scores; low happiness levels among the elderly with cognitive deficits have been associated with depression, disability, and loneliness (Tan et al., 2019). It has been demonstrated that identifying factors that influence the conversion of mild and moderate cognitive deficits to dementia can slow or prevent progressive deficits (Pan et al., 2021; Nelson et al., 2021; Silva et al., 2020; Sharma et al., 2020; Ma, 2020; Hu et al., 2021a). Based on the results of risk factor research, it seems possible to carry out preventive measures at an early stage to preserve and improve the cognitive functions of the aging population.

Kazakhstan is experiencing an accelerated rate of population aging (Vollset et al., 2020). The anticipated “epidemic of dementia” has prompted world-class experts to develop national plans to overcome its devastating socioeconomic impact (World Alzheimer’s Report, 2018). To date, a limited number of studies on dementia and risk factors in Central Asia and Kazakhstan have been conducted (Tsoy et al., 2019b; Madenbay et al., 2018; Manton et al., 2004; Askarova et al., 2020; Tsoy et al., 2019a). Determining the factors influencing the conversion of mild and moderate cognitive deficits to dementia disorders could help us to restrain or even prevent progressive deficits as moderate cognitive deficits constitute a transitional stage in developing dementia with a conversion range of 10-15% per year (Eshkoo et al., 2015). For this reason, it was decided to determine the socio-demographic factors and personality traits that predispose elderly patients in Kazakhstan with cognitive impairment non dementiatio the development of mixed anxiety-depressive disorders with cognitive deficits.

MATERIAL AND METHODS

During the research an observational study involving 111 patients from Kazakhstan with cognitive deficits was conducted: 59 patients, whose depressive disorders had been identified by clinical and psychopathological research using the Hamilton Depression Rating Scale (HRSD) and 52 patients for whom psychogenic depressive disorders had not been defined.

The recruitment of the study participants was carried out in the state nursing home for the elderly and disabled. The study included male and female individuals aged 60-74 years with different national and ethnic backgrounds, with mild to moderate cognitive impairment, who wished to participate in the study and signed an informed consent to participate in the study.

186 elderly people aged 60-74 years were examined. 111 patients were selected according to the leading selection criterion – the presence of cognitive impairment, not reaching the degree of severe dementia.

Criteria for exclusion of patients from the study: patients under 60 and over 75 years of age; patients with severe cognitive impairment due to the lack of the

possibility of purposeful communication (filling out informed consent, questionnaires) and performing test tasks related to this study; patients in an acute psychotic state; patients with a history of mental pathology (schizophrenia, affective disorders, traumatic brain damage, epilepsy, etc.); patients who refused to participate at any phase of the research.

The associative relationship of depression with three groups of predictors was assessed: personal characteristics (sex, age, marital status, education, attitude to religion, relationships with relatives, psychopathological burden of heredity, and the age of onset of cognitive deficits); premorbid personality pattern (isolation, dependence, insecurity, passivity, selfishness, hostility, a tendency to feel guilty, and a tendency for self-justification), and the type of personality accentuation, according to Leonhard-Schmieschek (demonstrativeness, stuckness, pedantry, excitability, hyperthymia, dysthymia, anxiety, exaltation, and emotiveness).

Microsoft Excel and the SPSS Statistics package (IBM) were used for statistical processing of the data. The standard method was used to calculate the specific gravity (%) and standard error of the mean (\pm SE). Pearson's chi-squared test and the odds ratio (OR) were used to measure the reliability and closeness of the relationship between depression and the predictors studied.

Patients were included in the research sample after getting acquainted with the information about the objectives and methods of the study and signing the informed consent. Cognitive impairment using the Mini Mental State Examination (MMSE), as well as subjective complaints of cognitive impairment (memory impairment, lack of concentration, etc.) were assessed.

Among 111 patients with cognitive impairment, a HRSD questionnaire, a survey, and a conversation to identify anxiety-depressive symptoms were conducted. All this was carried out by a psychiatrist, who had studied medical history of each patient. Additionally, the patients were examined by a neuropathologist and a clinical psychologist and a general practitioner.

Conversation and observation are the main clinical and psychopathological method for diagnosing mental disorders. Almost all patients had cerebrovascular and cardiovascular diseases, which indicates in favor of verification of vascular cognitive impairment.

At stage II, study and comparison of these groups according to the socio-demographic, personal-psychological, clinical and psychopathological data of the patients in the observed groups were occurred. It has been found out that two groups differ.

For instance, in the group with anxiety-depressive disorders, women, patients who did not have a spouse (died, divorced) predominated, and more often these patients had conflict relationships with close relatives. Also, in that group there were more of those patients who described themselves as private, uncommunicative, anxious and inactive individuals.

It can be assumed that these personality traits contributed to the fact that these patients were more often lonely, had conflict or had a cold relationship with relatives and given the fact that in Kazakhstan the homes of the elderly are quite

unpopular choice of destination, they were forced to go there. In Kazakhstan, the elderly is held in high esteem and usually they are less likely to live alone in old age.

Further, in this study there is a prescription of anti-dementia treatment for all patients (donepezil hydrochloride, memantine in dosages of either 5 or 10 mg according to the treatment protocol of Republic of Kazakhstan), also conducted cognitive training in both groups. In a group 1 – with anxiety-depressive disorders, additional antidepressants with anti-anxiety effects were prescribed and conducted psychotherapy. And the results of treatment were evaluated after 6 months, but this will be presented in another article. Below there is brief results description for the reference purpose.

RESULTS

An association between depression and sex ($p = 0.001$), marital status ($p \leq 0.001$), and the nature of the patients' relationship with relatives ($p=0.005$) was revealed. Further, it was found that the chance of developing depression was higher in female patients (OR=3.657), lonely patients (OR=5.984), and patients with conflicting relationships with relatives (OR=3.222). The frequency of depression was higher in patients with the personality traits of isolation (OR=2.624, $p=0.016$), insecurity (OR=2.879, $p=0.007$), and passivity (OR=2.175, $p=0.049$). Patients showing the "stuckness" effect (OR=2.924, $p=0.049$), pathological properties of dysthymia (OR = 2.279, $p = 0.035$), and anxiety (OR=4.345, $p=0.001$), were more likely to develop depression.

Following the diagnoses of depressive disorders in some patients using the HDRS, the existence of an associative relationship between depression and the patients' personal characteristics (common predictors) was assessed. It is clear now that depression was associated with sex ($\chi^2=10.845$, $p=0.001$), marital status ($\chi^2=13.708$, $p \leq 0.001$) and the nature of the patient's relationship with relatives ($\chi^2=8.068$, $p=0.005$) (Table 1).

Female had a 3.657-fold (in some cases, 8.020-fold) increased chance of a depressive disorder compared with male patients. The proportion of depressive disorders among women and men was $67.8 \pm 6.08\%$ and $36.5 \pm 6.68\%$, respectively.

Depressive disorders were 5.984-fold more common in single patients than in married patients (OR CI 2.180÷16.425). The proportion of single and married patients with depression was $63.1 \pm 5.27\%$ and $22.2 \pm 8.00\%$ respectively.

The incidence of depression was higher among patients with conflicting relationships with relatives than among patients with benevolent and (or) formal relationships: $70.7 \pm 7.11\%$ and $42.9 \pm 5.91\%$ respectively. The strength of the association between depression and the nature of the patient's relationship with relatives was high; OR=3.222 (CI 1.415÷7.335). The predominance of women in the group (CIND+Depressive disorder) – 67.8%, while in the comparison group (depressive disorder) the majority were men – 63.5%. This can be explained by the greater emotionality of women compared to men and confirms the literature

Table 1. Association of depression with patients' personal characteristics (common predictors)

Common predictors	Depression scale (according to the HDRS)				OR	χ^2 Pearson's, p-value	
	Depressive disorder group		Comparison group				
	Absolute number	%	Absolute number	%			
Gender						3,657 (1.667+8.020)	$\chi^2=10.845$, p = 0.001
female	40	67.8%	19	32.2%			
male	19	36.5%	33	63.5%			
Age						0,661 (0.304+1.435)	$\chi^2=1.101$, p = 0.294
over 65	34	49.3%	35	50.7%			
up to 65	25	59.5%	17	40.5%			
Marital status						5,984 (2.180+16.425)	$\chi^2=13.708$, p ≤ 0.001
single	53	63.1%	31	36.9%			
married	6	22.2%	21	77.8%			
Education level						1,051 (0.419+2.634)	$\chi^2=0.011$, p = 0.916
primary/secondary	47	53.4%	41	46.6%			
higher	12	52.2%	11	47.8%			
Religious beliefs						0,554 (0.222+1.384)	$\chi^2=1.622$, p = 0.203
atheist	10	41.7%	14	58.3%			
believer	49	56.3%	38	43.7%			
Relationship with relatives						3,222 (1.415+7.335)	$\chi^2=8.068$, p = 0.005
conflicting	29	70.7%	12	29.3%			
friendly and/or formal	30	42.9%	40	57.1%			
Psychopathological burden of heredity						1,986 (0.802+4.920)	$\chi^2=2.242$, p = 0.134
no	49	57.0%	37	43.0%			
yes	10	40.0%	15	60.0%			
Age of onset of cognitive deficits						0,869 (0.410+1.843)	$\chi^2=0.134$, p = 0.715
over 60	32	51.6%	30	48.4%			
up to 60	27	55.1%	22	44.9%			

that anxiety-depressive disorders in the general population are more common in women than in men.

Next the association between depression and the patients' premorbid personal characteristics was looked (Table 2). Depression was conditioned by the personality traits of isolation ($c^2=5.755$, $p=0.016$), insecurity ($c^2=7.360$, $p=0.007$), and passivity ($c^2=3.875$, $p=0.049$).

A higher proportion of depressive disorders were identified in patients displaying the premorbid personal characteristic of isolation than in patients who did not display this characteristic (67.4±7.15% versus 44.1±6.02%; OR 2.624, CI 1.182÷5.825).

Depressive disorders were more typical for patients with the premorbid personality characteristic of insecurity than for patients who did not have this personality characteristic (65.0±6.16% versus 39.2±6.84%; OR 2.879, CI 1.329÷ 6.235).

The incidence of depressive disorders was higher among passive patients than patients who did not display this characteristic (64.4±7.14% versus 45.5± 6.13%; OR 2.175, CI 0.998÷4.741). Thus, premorbid character traits like insecurity, passivity and isolation determined the increased vulnerability of elderly patients to stressful situations and increased the risk of developing psychogenic anxiety-depressive disorders.

Table 2. Association of depression with patients' premorbid personal characteristics

Premorbid personal characteristics	Depression scale (according to the HDRS)				OR	χ^2 Pearson's, p-value
	Depressive disorder+CIND		CIND			
	Absolute number	%	Absolute number	%		
Isolation						
yes	29	67.4%	14	32.6%	2.624 (1.182+5.825)	$\chi^2=5.755$, p=0.016
no	30	44.1%	38	55.9%		
Addiction						
yes	26	63.4%	15	36.6%	1.943 (0.882+4.282)	$\chi^2=2.749$, p=0.097
no	33	47.1%	37	52.9%		
Uncertainty						
yes	39	65.0%	21	35.0%	2.879 (1.329+6.235)	$\chi^2=7.360$, p=0.007
no	20	39.2%	31	60.8%		
Passivity						
yes	29	64.4%	16	35.6%	2.175 (0.998+4.741)	$\chi^2=3.875$, p=0.049
no	30	45.5%	36	54.5%		
Selfishness						
yes	19	38.8%	30	61.2%	0.348 (0.160+0.756)	$\chi^2=7.283$, p=0.007
no	40	64.5%	22	35.5%		
Hostility						
yes	13	52.0%	12	48.0%	0.942 (0.386+2.298)	$\chi^2=0.017$, p=0.896
no	46	53.5%	40	46.5%		
Tendency to feel guilty						
yes	34	60.7%	22	39.3%	1.855 (0.872+3.944)	$\chi^2=2.595$, p=0.107
no	25	45.5%	30	54.5%		
Tendency for self-justification						
yes	25	41.7%	35	58.3%	0.357 (0.164+0.776)	$\chi^2=6.919$, p=0.009
no	34	66.7%	17	33.3%		

Finally, the association between depressive disorders and character accentuations according to K. Leonhard and H. Schmieschek (Methodology for the study, 1970) was looked (Table 3). It was identified that depression was associated with the “stuckness” effect ($c^2=3.881$, $p=0.049$), and the pathological properties of dysthymic ($c^2=4.437$, $p=0.035$) and anxious personality types ($c^2=11.987$, $p=0.001$).

The incidence of depression was higher among patients showing a stuck type of personality accentuation than patients who did not show this effect (73.7±10.10% versus 48.9±5.21%; OR 2.924, CI 0.974÷8.785).

The proportions of depressive disorders in patients with and without dysthymic personality types were 64.6±6.90% and 44.4±6.26%, respectively (OR 2.279, CI 1.053÷4.936).

The incidence of depression was significantly higher in patients with an anxious personality type than in patients who did not express this personality type (75.0±6.85% versus 40.8±5.83%; OR 4.345, CI 1.842÷10.247). The worsening of premorbid character traits such as isolation, passivity, uncertainty in elderly patients against the background of the development of vascular cognitive impairment contributed to the formation of accentuations (strengthening) of character traits mainly of the type of “anxiety”, “dysthymia” and a tendency to “get stuck” on emotionally negatively colored experiences, which made these patients vulnerable

Table 3. Association of depression with patients' personality accentuations according to K. Leonhard and H. Schmieschek

Personality accentuations according to K. Leonhard and H. Schmieschek	Depression scale (according to the HDRS)				OR	χ^2 Pearson's, p-value
	Depressive disorder+CIND		CIND			
	Absolute number	%	Absolute number	%		
Demonstrativeness						
tendency and/or accentuation	10	33.3%	20	66.7%	0.327 (0.135+0.787)	$\chi^2=6.486$, p=0.011
not expressed	49	60.5%	32	39.5%		
Stuckness						
tendency and/or accentuation	14	73.7%	5	26.3%	2.924 (0.974+8.785)	$\chi^2=3.881$, p=0.049
not expressed	45	48.9%	47	51.1%		
Pedantry						
tendency and/or accentuation	12	63.2%	7	36.8%	1.641 (0.593+4.542)	$\chi^2=0.921$, p=0.337
not expressed	47	51.1%	45	48.9%		
Excitability						
tendency and/or accentuation	17	37.0%	29	63.0%	0.321 (0.146+0.704)	$\chi^2=8.276$, p=0.004
not expressed	42	64.6%	23	35.4%		
Hyperthymia						
tendency and/or accentuation	17	43.6%	22	56.4%	0.552 (0.251+1.213)	$\chi^2=2.208$, p=0.137
not expressed	42	58.3%	30	41.7%		
Dysthymia						
tendency and/or accentuation	31	64.6%	17	35.4%	2.279 (1.053+4.936)	$\chi^2=4.437$, p=0.035
not expressed	28	44.4%	35	55.6%		
Anxiety						
tendency and/or accentuation	30	75.0%	10	25.0%	4.345 (1.842+10.247)	$\chi^2=11.987$, p=0.001
not expressed	29	40.8%	42	59.2%		
Exaltation						
tendency and/or accentuation	8	29.6%	19	70.4%	0.272 (0.107+0.694)	$\chi^2=7.929$, p=0.005
not expressed	51	60.7%	33	39.3%		
Emotiveness						
tendency and/or accentuation	27	42.9%	36	57.1%	0.375 (0.172+0.818)	$\chi^2=6.202$, p=0.013
not expressed	32	66.7%	16	33.3%		
Cyclothymia						
tendency and/or accentuation	15	50.0%	15	50.0%	0.841 (0.364+1.945)	$\chi^2=0.164$, p=0.685
not expressed	44	54.3%	37	45.7%		

to stressful situations and contributed to the development of psychogenic anxiety-depressive disorders.

The mean value of the Hamilton Depression Scale (HDRS, HAM-D) before the complex intervention was 9.8 (standard deviation 1.5), after the interventions the mean value was 3.7 (SD±1.0). It can be qualified as a complete reduction of depressive disorders. The values of the difference in the test results disprove the null hypothesis, and confirm that, the means in the two groups after the interventions are unequal, and the mean difference is 6.119 (SD±1.566), CI (5.711; 6.527), p -value=0.000.

A similar positive dynamics (during treatment) therapy was observed in the level of anxiety. The mean score of the level of anxiety on the Hamilton anxiety scale (HDRS, HAM-Anxiety) before treatment was 15.0 points (SD±2.7), and after therapy this the indicator dropped to 5.1 points (SD±1.3). The test data in-

indicated that the null hypothesis should be refuted, since the average difference between the indicators was 9.881 (SD±2.738), CI (9.174; 10.589), p -value= 0.000.

It was assessed not only the dynamics of anxiety and depressive disorders, but also the level of cognitive impairment after 6 months of observation. In the group of patients with VCI, before treatment, the mean score of cognitive functions according to the MMSE Scale was 25.9 points (SD±1.2), after treatment the mean score result rose up to 26.7 points (SD±0.7). The difference in mean score before and after treatment was 0.8 points (SD±1.09), CI 1.074; -0.464, and p -value 0.000. In this case, it was managed to suspend the development of cognitive impairment.

In the group of patients with VCI + ADD (anxiety-depression disorder), the mean score of cognitive functions according to the MMSE Scale was 24.4 (SD ±0.6), after treatment the mean score result was 27.2 points (SD±1.0). The difference in values before and after treatment was 2.79 (SD±1.229), CI -3.117; -2.476, and p -value 0.000.

In the group, good treatment results were observed, given that an additionally application of psychopharmacotherapy and psychotherapeutic methods to treat psychotraumatic factors and personality traits, which were assumed to worsen the patients' conditions.

In Kazakhstan, the problem is that elderly patients often complain about the deterioration of cognitive activity, but doctors do not examine the problem comprehensively. Causes are not fully explored. Often, could be observed the situation when an older woman (widowed and abandoned) having deteriorated memory comes to see the doctor, while being already inuntreated stage.

And it often happens in the clinic that the grandmother's memory has completely deteriorated after the grandfather died, or she was left alone, the children all parted. And see a doctor in later stages.

DISCUSSION

Many researchers have investigated how various factors influence the development of mixed anxiety-depressive disorders in the elderly population (Hellwig and Domschke, 2019; Suradom et al., 2019; Beghi et al., 2021; Van der Heide et al., 2021; Alexopoulos, 2019). However, few studies have focused on determining the factors that might mediate the development of mixed anxiety-depressive disorders among the elderly with pre-existing cognitive disorders of a non-dementia level. Due to a problematic of this knowledge gap, attention of the researchers was focused on identifying the socio-psychological characteristics that might mediate the development of mixed anxiety-depressive disorders in elderly patients with cognitive deficits of a non-dementia level.

Research results show that in elderly patients with anxiety-depressive disorders, the loss of a spouse, breakdown of family relations, and unfavorable, especially conflicting, relationships with other relatives strong predictors of depressive disorder development in elderly patient with cognitive impairment non dementia. Notably that the single patients in the cohort were nearly six times more likely to

have depressive disorders than married patients. It also found that patients who had conflicting relationships with close relatives had a three-fold higher risk of developing depression than patients with benevolent and/or formal relationships. Previous studies have identified that elderly people in social isolation, living alone, and with health stressors as having more frequent occurrences of depression (Sinyor et al., 2016). The latter is possibly a protective factor; however, the role of protective factors was not investigated in this study and this would be an interesting area of future research.

The nature of relationships with loved ones is the most significant factor in the development of anxiety and depression. More elderly people may be alone nowadays as the extended family model, in which an elderly person remains in demand for a long time and is not alone, has gradually lost its relevance. Many have reported that social isolation, bereavement, psychological pain, lack of significant social contacts, a sense of one's own invisibility, meaninglessness of existence, cumulative suffering, and a "painful life" contribute to increased suicidal activity among the elderly (Wand et al., 2018; Conejero et al., 2018; Pilania et al., 2019; Conejero et al., 2016; Shah et al., 2016).

While many have investigated the effects of psycho-traumatic factors on dementia, few have examined the personality traits that determine increased sensitivity to stressors (Mosazadeh et al. 2023). An examining the aspects of character and temperament in the elderly cohort, who already show cognitive deficits, was in focus. In those with mixed anxiety-depressive disorders, those with the personality accentuations of anxiety, dysthymia, and "stuckness" were XX times more likely to have experienced negative emotional experiences than those in without anxiety-depressive disorders. The premorbid personality traits of passivity, insecurity, and isolation also prevailed in this group of elderly patients with mixed anxiety-depressive disorders. All of these personality traits determined the increased vulnerability of these patients to stressful situations and increased the risk of developing psychogenic mixed anxiety-depressive disorders. Others have consistently reported that patients with higher rates of anxiety, negative emotions, hostility, pessimism, hopelessness, and susceptibility to limitations had a 20–30% increased risk of dementia and a 10–20% increased risk of cognitive impairment no dementia (CIND) (Sutin et al., 2018a). Higher susceptibility to dementia in patients with a high neurotic score has been reported in meta-analyses (Terracciano et al., 2014).

In Kazakhstan, elderly people with cognitive deficits receive antedementia therapy typically in the form of donepezil (and other cholinesterase inhibitors) and memantine; however, this treatment does not affect the social and psychological factors that worsen the cognitive deficits. With a more thorough examination of the patient, including their socio-psychological status, it might be possible to provide more comprehensive assistance and reduce the risk of mild cognitive impairment converting to dementia.

CONCLUSIONS

During this research the main reasons of the likelihood of patients with cognitive deficits developing a depressive disorder were found. Among such medical features could be listed patients' personal characteristics (gender, marital status, and the nature of the patients' relationship with relatives), premorbid personality patterns (isolation, insecurity, and passivity), and the pathological personality properties of stuckness, dysthymia and anxiety.

The study was based on the observation and analysis of data from old persons, who were selected according to specific characteristics and divided into two groups. A six-months study of the preconditions for depression in old age revealed that women are more likely to develop such diseases in old age than men are.

In summary, the analysis of the clinical features of elderly patients with cognitive deficits suggests that their likelihood of developing mixed anxiety-depressive disorders is, to a certain extent, determined by their personal characteristics, premorbid personality, pathological personality traits and negative emotional experiences. While many of findings in this research are supported by data conducted in other cohorts, the study itself is one of the first in this medical area, and there was no similar research conducted previously in Kazakhstan. Going forward, it is desirable to implement a greater coverage of the individuals studied to create diagnostic tools to identify risk groups among the elderly population in Kazakhstan. In the Republic of Kazakhstan, the elderly not only from the nursing home, but also look at the groups of the elderly who live with their families could be covered. In general, it could be an effective step for prevention such disease in a way of introducing some criteria for risk groups among the elderly. Current situation in Kazakh hospitals: if you are 40 and you are a woman, then you will definitely be screened for breast cancer. What can be implemented in the future: if you are 60 you are taking a test for MILD cognitive impairment, if you have lost your spouse / live alone / you have a conflict relationship in the family, then the identification of anxiety-depressive symptoms (this can be a short 5-minute screening questionnaire). There is a strong belief that in doing so, it will not only be possible to more accurately diagnose and treat affected individuals, but also establish preventive measures for the elderly with cognitive and mixed anxiety-depressive disorders.

REFERENCES

- Alexopoulos, G.S. 2019. Mechanisms and treatment of late-life depression. *Transnational Psychiatry*, 9(1), 188.
- Askarova, S., Umbayev, B., Masoud, A.-R., Kaiyrylkyzy, A., Safarova, Y., Tsoy, A., Olzhayev, F., Kushugulova, A. 2020. The links between the gut microbiome, aging, modern lifestyle and Alzheimer's disease. *Frontiers in Cellular and Infection Microbiology*, 10, 104.
- Beghi, M., Butera, E., Cerri, C.G., Cornaggia, C.M., Febbo, F., Mollica, A., Berardino, G., Piscitelli, D., Resta, E., Logroscino, G., Daniele, A., Altamura, M., Bellomo, A., Panza, F., Lozupone, M. 2021. Suicidal behaviour in older age: a systematic review of risk factors associated to suicide attempts and completed suicides. *Neuroscience & Biobehavioral Review*, 127, 193-211.

- Conejero, I., Lopez-Castroman, J., Giner, L., Baca-Garcia, E. 2016. Sociodemographic antecedent validators of suicidal behavior: a review of recent literature. *Current Psychiatry Reports*, 18(10), 94.
- Conejero, I., Olié, E., Courtet, P., Calati, R. 2018. Suicide in older adults: current perspectives. *Clinical Interventions in Aging*, 13, 691-699.
- Delfino, L.L., Komatsu, R.S., Komatsu, C., Neri, A.L., Cachioni, M. 2018. Path analysis of caregiver characteristics and neuropsychiatric symptoms in Alzheimer's disease patients. *Geriatrics & Gerontology International*, 18(8), 1177-1182.
- Eshkooor, S.A., Hamid, T.A., Mun, C.Y., Chee, K.Ng. 2015. Mild cognitive deficits and its management in older people. *Clinical Interventions in Aging*, 10(10), 687-93.
- Fischer, C.E., Kotrebi, I., Karameh, W.K., Kumar, S., Gallagher, D., Golas, A., Munoz, D., Barfett, H., Butters, M.A., Bowie, Ch.R., Flint, A., Rajji, T., Herrmann, N., Pollock, B.G., Mulsant, B., Schweizer, T.A., Mah, L., the PACT-MD Study Group. 2019. Examining the link between cardiovascular risk factors and neuropsychiatric symptoms in mild cognitive impairment and major depressive disorder in remission. *Journal of Alzheimer's Disease*, 67(4), 1305-1311.
- Floud, S., Balkwill, A., Sweetland, S., Brown, A., Reus, E.M., Hofman, P.A. 2021. Cognitive and social activities and long-term dementia risk: the prospective UK Million Women Study. *The Lancet. Public Health*, 6(2), e116-e123.
- Fu, J., Jiang, Z., Hong, Y., Liu, S., Kong, D., Zhong, Zh., Luo, Y. 2021. Global scientific research on social participation of older people from 2000 to 2019: A bibliometric analysis. *International Journal of Older People Nursing*, 16(1), e12349.
- Gigi, A., Papirovit, M. 2021. Association of anxiety awareness with risk factors of cognitive decline in MCI. *Brain Sciences*, 11(2), 135.
- Ginsberg, T.B., Powell, L., Emrani, Sh., Wasserman, V., Higgins, S., Chopra, A., Cavalieri, T.A., Libon, D.J. 2019. Instrumental activities of daily living, neuropsychiatric symptoms, and neuropsychological impairment in mild cognitive deficits. *The Journal of the American Osteopathic Association*, 119(2), 96-101.
- Hellwig, S., Domschke, K. 2019. Anxiety in late life: an update on pathomechanisms. *Gerontology*, 65(5), 465-473.
- Hu, C., Wang, L., Zhao, X., Zhu, B., Tian, M., Qin, H. 2021. Investigation of risk factors for the conversion of mild cognitive deficits to dementia. *The International Journal of Neuroscience*, 131(12), 1-8.
- Hu, C.P., Zhao, Y.-H., Zhao, X.-D., Zhu, B.-G., Qin, H.Y. 2021. Correlation between sleep characteristics and cognitive decline in the elderly people: A cross-sectional study in China. *International Journal of Clinical Practice*, 75(7), e14225.
- Ikezaki, H., Hashimoto, M., Ishikawa, T., Fukuhara, R., Tanaka, H., Yuki, S., Kuribayashi, K., Hotta, M., Koyama, A., Ikeda, M., Takebayashi, M. 2020. Relationship between executive dysfunction and neuropsychiatric symptoms and impaired instrumental activities of daily living among patients with very mild Alzheimer's disease. *International Journal of Geriatric Psychiatry*, 35(8), 877-887.
- Ilesheva, R.G. 1981. *Affective psychosis in later life*. Alma-Ata: Nauka, 135 p.
- Izmailova, N.T., Ilesheva, R.G., Kudyarova, G.M. 2000. Features of the age aspect of depressive disorders. In: *Topical issues of psychiatry, narcology and psychology: Collection of scientific (pp. 69-72)*. Almaty: Nauka.
- Jenkins, A., Tree, J., Tales, A. 2021. Distinct profile differences in subjective cognitive decline in the general public are associated with metacognition, negative affective symptoms, neuroticism, stress, and poor quality of life. *Journal of Alzheimer's Disease*, 80(3), 1231-1242.
- Jia, L., Du, P.Y., Chu, P.L., Zhang, P.Zh., Li, F., Lyu, D., Li, Y., Zhu, M., Jiao, H., Song, Y., Shi, Y., Zhang, H., Gong, M., Wei, C., Tang, Y., Famng, P.B., Guo, D., Wang, F., Zhou, A., Chu, Ch., Zuo, X., Yu, Y., Yuan, Q., Wang, W., Li, P.F., Shi, P.Sh., Yang, H., Zhou, P.Ch., Liao, Zh., Lv, Y., Li, Y., Kan, M., Zhao, P.H., Wang, Sh., Yang, Sh., Li, H., Liu, Zh., Wang, Q., Qin, W., Hia, P.J. 2020. Prevalence, risk factors, and management of dementia and mild cognitive impairment in adults aged 60 years or older in China: a cross-sectional study. *The Lancet. Public Health*, 5(12), e661-e671.
- Johansson, L., Guo, X., Duberstein, P.R., Hällström, T., Waern, M., Ostling, S., Skoog, I. 2014. Midlife personality and risk of Alzheimer disease and distress: a 38-year follow-up. *Neurology*, 83(17), 1538-1544.

- Kim, H.J., Lee, J.-H., Cheong, E.-N., Chung, S.-E., Jo, S., Shim, W.-H., Jomg, Y.-J. 2020. Elucidating the risk factors for progression from amyloid-negative amnesic mild cognitive defects to dementia. *Current Alzheimer Research*, 17(10), 893-903.
- Krivanek, T.J., Gale, S.A., McFreeley, B.M., Nicastrì, C.M., Daffner, K.R. 2021. Promoting successful cognitive aging: a ten-year update. *Journal of Alzheimer's Disease*, 81(3), 871-920.
- Kudyarova, G.M. 1989. Depressions of involuntary age (2nd ed.). Alma-Ata: Nauka, 168p.
- Lacour, A., Espinosa, A., Louwersheimer, E., Heilmann, S., Hernández, I., Wolfsgruber, S., Fernández, V., Wagner, H., Rosende-Roca, M., Mauleón, A., Moreno-Grau, S., Vargas, L., Pijnenburg, Y.A.L., Koene, T., Rodríguez-Gómez, O., Ortega, G., Ruiz, S., Holstege, H., Sotolongo-Grau, O., Kornhuber, J., Peters, O., Frölich, L., Hüll, M., Rütther, E., Wiltfang, J., Scherer, M., Riedel-Heller, S., Alegret, M., Nöthen, M.M., Scheltens, P., Wagner, M., Tárraga, L., Jessen, F., Boada, M., Maier, W., van der Flier, W.M., Becker, T., Ramirez, A., Ruiz, A. 2017. Genome-wide significant risk factors for Alzheimer's disease: role in progression to dementia due to Alzheimer's disease among subjects with mild cognitive defects. *Molecular Psychiatry*, 22(1), 153-160.
- Lara, E., Koyanagi, A.I., Domènech-Abella, J., Miret, M., Ayuso-Mateos, J.L., Haro, J.M. 2017. The impact of depression on the development of mild cognitive defects over 3 years of follow-up: a population-based study. *Dementia and Geriatric Cognitive Disorders*, 43(3-4), 155-169.
- Ma, L. 2020. Depression, anxiety, and apathy in mild cognitive defects: current perspectives. *Frontiers in Aging Neuroscience*, 12, 9.
- Madenbay, K., Shalkharova, Zh., Nuskabayeva, G., Sadykova, K. 2018. Association between components of metabolic syndrome and cognitive dysfunction: cross-sectional study among the population of the Turkestan region. *Georgian Medical News*, 278, 114-120.
- Manton, K.G., Volovik, S., Kulminski, A. 2004. ROS effects on neurodegeneration in Alzheimer's disease and related disorders: on environmental stresses of ionizing radiation. *Current Alzheimer Research*, 1(4), 277-293.
- McGrattan, A.M., Zhu, Y., Richardson, C.D., Mohan, D., Soh, Y.Ch., Sajjad, A., van Aller, C., Chen, Sh., Paddick, S.-M., Prina, M., Siervo, M., Robinson, L.A., Stephan, B.C.M. 2021. Prevalence and risk of mild cognitive defects in low and middle-income countries: a systematic review. *Journal of Alzheimer's Disease*, 79(2), 743-762.
- Methodology for the study of personality accentuations. Stuck type of human nature. 1970. <https://basheschool2.ru/en/oprosnik-leongarda-shmisheka-opisanie-metodika-izucheniya-akcentuacii.html>
- Mora, F. 2013. Successful brain aging: plasticity, environmental enrichment, and lifestyle. *Dialogues in Clinical Neuroscience*, 15(1), 45-52.
- Mosazadeh H., Davoodi P., Rajabi A., Blachnio A. (2023). COVID-19 fear and Health-Related Quality of Life of older adults: evidence from Iran. *Acta Neuropsychologica* 2023; 21 (1): 21-31. DOI: 10.5604/01.3001.0016.3034.
- Nelson, M.E., Jester, D.J., Petkus, A.J., Andel, R. 2021. Cognitive reserve, Alzheimer's neuropathology, and risk of dementia: a systematic review and meta-analysis. *Neuropsychology review*, 31(2), 233-250.
- Nolte, E.D., Nolte, K.A., Yan, S.S. 2019. Anxiety and task performance changes in an aging mouse model. *Biochemical and Biophysical Research Communications*, 514(1), 246-251.
- Pais, R., Ruano, L., Moreira, C., Carvalho, O.P., Barros, H. 2020. Prevalence and incidence of cognitive impairment in an elder Portuguese population (65-85 years old). *BMC Geriatrics*, 20(1), 470.
- Pan, Y., Shea, Y.-F., Li, S., Chen, R., Mak, H.K.-F., Chiu, P.K.-Ch., Chu, L.-W., Song, Y.-Q. 2021. Prevalence of mild behavioural impairment: a systematic review and meta-analysis. *Psychogeriatrics: the Official Journal of the Japanese Psychogeriatric Society*, 21(1), 100-111.
- Petkus, A.J., Reynolds, Ch.A., Wetherell, J.L., Kremen, W.S., Gatz, M. 2017. Temporal dynamics of cognitive performance and anxiety across older adulthood. *Psychology and Aging*, 32(3), 278-292.
- Pilania, M., Yadav, V., Bairwa, M., Behera, P., Gupta, S.D., Khurana, H., Mohan, V., Baniya, G., Poongothai, S. 2019. Prevalence of depression among the elderly (60 years and above) po-

- pulation in India, 1997-2016: a systematic review and meta-analysis. *BMC Public Health*, 19(1), 832.
- Rainero, I., Bruni, A.C., Marra, C., Cagnin, A., Bonanni, L., Cupidi, Ch., Lagana, V., Rubino, E., Vacca, A., Lorenzo, R.D., Provero, P., Isella, V., Vanacore, N., Agosta, F., Appollonio, I., Caffarra, P., Busse, C., Sambati, R., Quaranta, D., Guglielmi, V., Logroscino, G., Filippi, M., Tedeschi, G., Ferrarese, C., SINdem COVID-19 Study Group. 2020. The impact of COVID-19 quarantine on patients with dementia and family caregivers: a nation-wide survey. *Frontiers in Aging Neuroscience*, 12, 625781.
- Romero-Sevilla, R., Casado-Naranjo, I., Portilla-Cuenca, J.C., San Juan, B.D., Fuentes, J.M., Lopez-Espuela, F. 2018. Vascular risk factors and lesions of vascular nature in magnetic resonance as predictors of progression to dementia in patients with mild cognitive defects. *Current Alzheimer Research*, 15(7), 671-678.
- Seifan, A., Ganzer, Ch.A., Ryon, K., Lin, M., Mahmudur, R., Adolfo, H., Shih, C., Jacobs, A.R., Greenwald, M., Isaacson, R.S. 2019. Detecting non-cognitive features of prodromal neurodegenerative diseases. *Current Aging Science*, 11(4), 242-249.
- Shah, A., Bhat, R., Zarate-Escudero, S., DeLeo, D., Erlangsen, A. 2016. Suicide rates in five-year age-bands after the age of 60 years: the international landscape. *Aging & Mental Health*, 20(2), 131-8.
- Sharma, G., Parihar, A., Talaiya, T., Dubey, K., Porwal, B., Parihar, M.S. 2020. *cognitive defects in type 2 diabetes, risk factors and preventive strategies*. *Journal of basic and clinical physiology and pharmacology*, 31(2). DOI: 10.1515/jbcpp-2019-0105
- Silva, D., Cardoso, S., Guerreiro, M., Maroco, J., Mendes, T., Alves, L., Nogueira, J., Baldeiras, I., Santana, I. 2020. Neuropsychological contribution to predict conversion to dementia in patients with mild cognitive defects due to Alzheimer's disease. *Journal of Alzheimer's Disease*, 74(3), 785-796.
- Sinyor, M., Tan, L.P., Schaffer, A., Gallagher, D., Shulman, K. 2016. Suicide in the oldest old: an observational study and cluster analysis. *International Journal of Geriatric Psychiatry*, 31(1), 33-40.
- Suradom, C., Wongpakaran, N., Wongpakaran, T., Lerttrakarnnon, P., Jiraniramai, S., Taemeeyapradit, U., Lertkachatarn, S., Arunpongpaisal, S. 2019. Prevalence and associated factors of comorbid anxiety disorders in late-life depression: findings from geriatric tertiary outpatient settings. *Neuropsychiatric Diseases and Treatment*, 15, 199-204.
- Sutin, A.R., Luchetti, M., Stephan, Y., Terracciano, A. 2021. Purpose in life and motoric cognitive risk syndrome: replicable evidence from two national samples. *Journal of the American Geriatrics Society*, 69(2), 381-388.
- Sutin, A.R., Stephan, Y. Terracciano, A. 2018. Psychological distress, self-beliefs, and risk of cognitive defects and dementia. *Journal of Alzheimers Disease*, 65(3), 1041-1050.
- Sutin, A.R., Stephan, Y., Terracciano, A. 2018. Psychological well-being and risk of dementia. *International Journal of Geriatric Psychiatry*, 33(5), 743-747.
- Tan, J.H., Abdin, E., Shahwan, S., Zhang, Y., Sambasivam, R., Vaingankar, J.A., Mahendran, R., Chua, H.C., Chong, S.A., Subramaniam, M. 2019. Happiness and cognitive defects among older adults: investigating the mediational roles of disability, depression, social contact frequency, and loneliness. *International Journal of Environmental Resaerch and Public Health*, 16(24), 4954.
- Terracciano, A., Sutin, A.R., An, Y., O'Brien, R.J., Ferrucci, L., Zonderman, A. B., Resnick, S.M. 2014. Personality and risk of Alzheimer's disease: new data and meta-analysis. *Alzheimer`s & Dementia*, 10(2), 179-186.
- Thancharoen, O., Waleekhachonloet, O., Limwattananon, Ch., Anutrakulchai, S. 2020. Cognitive impairment, quality of life and healthcare utilization in patients with chronic kidney disease stages 3 to 5. *Nephrology*, 25(8), 625-633.
- Tsoy, A., Saliev, T., Abzhanova, E., Turgambayeva, A., Kaiyrykzy, A., Akishev, M., Saparbayev, S., Umbayev, B., Askarova, Sh. 2019. The effects of mobile phone radiofrequency electromagnetic fields on beta-amyloid-induced oxidative stress in human and rat primary astrocytes. *Neuroscience*, 408, 46-57.

- Tsoy, R.T., Turuspekova, S.T., Klipitskaya, N.K., Mereke, A., Cumming, R.G. 2019. Prevalence of mild cognitive defects among older people in Kazakhstan and potential risk factors: across-sectional study. *Alzheimer Disease and Associated Disorders*, 33(2), 136-141.
- Van der Heide, F.C.T., Steens, I.L.M., Geraets, A.F.J., Foreman, Y.D., Henry, R.M.A., Kroon, A.A., van der Kallen, C.J.H., van Sloten, T.T., Dagnelie, P.C., van Dongen, M.C.J.M., Eussen, S.J.P.M., Berendschot, T.T.J.M., Schouten, J.S.A.G., Webers, C.A.B., van Greevenbroek, M.M.J., Wesselius, A., Koster, A., Schaper, N.C., Schram, M.T., Köhler, S., Stehouwer, C.D.A. 2021. Association of retinal nerve fiber layer thickness, an index of neurodegeneration, with depressive symptoms over time. *JAMA Network Open*, 4(11), e2134753.
- Vicini Chilovi, B., Conti, M., Zanetti, M., Mazzù, I., Rozzini, L., Padovani, A. 2009. Differential impact of apathy and depression in the development of dementia in mild cognitive defects patients. *Dementia and Geriatric Cognitive Disorders*, 27(4), 390-398.
- Vollset, S.E., Goren, E., Yuan, Ch.-W., Cao, J., Smith, A.E., Hsiao, T., Bisignano, C., Azhar, G.S., Castro, E., Chalek, J., Dolgert, A.J., Frank, T., Fukutaki, K., Hay, S.I., Lozano, R., Mokdad, A.H., Nandakumar, V., Pierce, M., Pletcher, M., Robalik, T., Steuben, K.M., Wunrow, H.Y., Zlavog, B.S., Murray, Ch.J.L. 2020. Fertility, mortality, migration, and population scenarios for 195 countries and territories from 2017 to 2100: a forecasting analysis for the Global Burden of Disease Study. *Global Health Metrics*, 396, 1285-1306.
- Wand, A.P.F., Peisah, C., Draper, B., Brodaty, H. 2018. Understanding self-harm in older people: a systematic review of qualitative studies. *Aging and Mental Health*. 22(3), 289-298.
- Watanabe, Y., Raaphorst, J., Izumi, Y., Yoshino, H., Ito, S., Adachi, S., Takigawa, H., Masuda, M., Atsuta, N., Adachi, Y., Iose, S., Arai, K., Yokota, O., Oda, M., Ogino, M., Ichikawa, H., Hasegawa, K., Kimura, H., Shimizu, T., Aiba, I., Yabe, H., Kanba, M., Kusumi, K., Aoki, T., Hiroe, Y., Watanabe, H., Nishiyama, K., Nomoto, M., Sobue, G., Beeldman, E., Hanajima, R., Nakashima, K., ALS_FTD_Q_J research group. 2020. Cognitive and behavioral status in Japanese ALS patients: a multicenter study. *Journal of Neurology*, 267(5), 1321-1330.
- Wiegelmann, H., Speller, S., Verhaert, L.-M., Schirra-Weirich, L., Wolf-Ostermann, K. 2021. Psychosocial interventions to support the mental health of informal caregivers of persons living with dementia - a systematic literature review. *BMC Geriatrics*, 21(1), 94.
- World Alzheimer's Report. 2018. <https://www.alz.co.uk/research/world-report-2018>
- World Population Ageing. 2017. https://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA_2017_Highlights.pdf
- Yang, A.N., Wang, X.-L., Rui, H.-R., Luo, H., Pang, M., Dou, X.-M. 2020. Neuropsychiatric symptoms and risk factors in mild cognitive defects: a cohort investigation of elderly patients. *The Journal of Nutrition, Health & Aging*, 24(2), 237-241.
- Yatawara, C., Hiu, Sh., Tan, L., Kandiah, N. 2018. Neuropsychiatric symptoms in South-East Asian patients with mild cognitive impairment and dementia: prevalence, subtypes, and risk factors. *International Journal of Geriatric Psychiatry*, 33(1), 122-130.
- Zaltsman, I.G. 1998. Diagnosis of depressive states. Almaty: Nauka, 37p.
- Zufferey, V., Gunten, A.V., Kherif, F. 2020. Interactions between personality, depression, anxiety and cognition to understand early stage of Alzheimer's disease. *Current Topics in Medicinal Chemistry*, 20(9), 782-791.

Corresponding author

Madina Kassymzhanova

Department of Psychiatry and Narcology

Kazakh National Medical University named after S.D. Asfendiyarov

050000, 94 Tole bi Str., Almaty, Republic of Kazakhstan

e-mail: madinakassymzhanovapsy@gmail.com