

EXERCISE TRAINING-INDUCED CHANGES IN INFLAMMATORY MEDIATORS AND HEAT SHOCK PROTEINS IN CANOEISTS

Barbara Morawin,^{1, C, D} Mateusz Rynkiewicz,^{2, A, B, E} Joanna Orysiak,^{3, C, D}
Tadeusz Rynkiewicz,^{1, A, B, E} Agnieszka Zembroń-Łacny^{4, A, B, E}

¹ University School of Physical Education, Poznań, Poland

² Diagnostic Sport Centrum Fit-Test, Gorzów Wielkopolski, Poland

³ Department of Nutrition Physiology, Institute of Sport, Warsaw, Poland

⁴ Department of Biology Basis of Physical Education and Sport, University of Zielona Gora, Poland

^A Study Design; ^B Data Collection; ^C Statistical Analysis; ^D Manuscript Preparation; ^E Funds Collection

The study was carried out at University School of Physical Education Poznan.

Address for correspondence:

Barbara Morawin, MSc

University School of Physical Education Poznan; Doctoral Study

Królowej Jadwigi 27/39, 61-871 Poznań, Poland

Phone: 667145947

E-mail: barbara.morawin@gmail.com

Abstract. According to cytokine overtraining theory, skeletal muscle injuries are related to systemic inflammatory reaction. In response to inflammation, cells rapidly produce a series of proteins known as heat shock proteins (HSPs). These are considered to be molecular chaperones which play a universal role in maintaining cellular homeostasis. Among the subset of stress-responsive proteins, HSP27 and HSP70 are considered to be a new approach to monitoring exercise training and adaptive mechanisms. The study was designed to demonstrate the effect of sport training on changes in pro-inflammatory cytokines and HSPs, and their relation with muscle damage and body composition.

Six elite canoeists (19.8 ± 2.9 yr) were observed during preparatory training period (March) at the 1st, the 4th and after 7 days of the conditioning camp, and then after 3 days of recovery. The canoeing training did not induce muscle damage, decreased in IL-1β and HSP27, increased in TNFα and HSP70 concentrations. The highest changes in TNFα and HSP70 were observed 3 days after conditioning camp (during recovery) compared to initial level (the 1st day of conditioning camp). TNFα correlated with HSP27 ($r = -0.563$; $P < 0.01$) and HSP70 ($r = 0.651$; $P < 0.001$). Any significant changes in body composition were not observed.

In conclusion, we could say that typical canoeing training improves cytokines and HSPs release, however, the changes are not related to muscle damage.

Key words: inflammation, cytokines, HSP27, HSP70, muscle damage

Introduction

The effectiveness of physical training depends on the training load, the individual toleration ability and the imbalance between the two may lead to under or over-training. One of the unique features of an exercise is that

it leads to a simultaneous increase of antagonistic mediators. On the one hand, exercise elevates catabolic pro-inflammatory cytokines such as interleukin 1 β (IL-1 β) and tumour necrosis factor α (TNF α). On the other hand, exercise stimulates anabolic components such as heat shock proteins (HSPs) which protect against stressors. If the anabolic response is stronger, exercise will probably lead ultimately to an increased muscle mass and exercise adaptation (Noble et al. 2008; Pedersen 2011; Roubenoff 2007).

The role of pro-inflammatory cytokines in skeletal muscle growth still has not been fully explored. It has been observed that after IL-1 β stimulation the total of protein synthesized does not increase, but rather the synthesis of the acute phase proteins is favoured (Weissman 1990). A study by Tayek (1996) has shown that TNF α has significant short- and long-term effects on protein synthesis. Earlier work demonstrated that TNF α can reduce weight gain and enhance muscle catabolism (Tracey et al. 1988). Nevertheless, the suppression of TNF α synthesis by using anti-inflammatory drug delays muscle restoration, but an excessive IL-1 β and TNF α release may be responsible for the overtraining (Mackey et al. 2007; Main et al. 2009). The measurement of both pro-inflammatory cytokines IL-1 β and TNF α within a population of athletes during training has not been widely reported (Borges et al. 2013; Main et al. 2009; Marin et al. 2011; Reinke et al. 2009; Zembron-Lacny et al. 2010). Nowadays, it is known that pro-inflammatory cytokines are elevated in sport activity and can be markers of overtraining syndrome, but we still have no information which level of inflammatory mediators is appropriate for athletes.

Heat shock proteins (HSPs) represent cell-protective system that may be induced by reactive oxygen species, cytokines, and hyperthermia. Under physiological conditions, constitutively expressed HSPs function as molecular chaperones, whereas under stress conditions, HSPs protect proteins against misfolding, aggregation and denaturation. In addition, HSPs may directly regulate specific stress- responsive signalling pathways and may antagonize signalling cascades that result in apoptosis (Madamanchi et al. 2001; Noble et al. 2008). Exercise-induced muscle damage is considered to be one of the stimulus which induce HSPs (Steinacker et al. 2004). Among the subset of stress-responsive proteins, HSP27 and HSP70 are considered as a new approach of monitoring exercise training and adaptive mechanisms (Banfi et al. 2006). The fibre type-specific expression of HSP70 is influenced by resistance and endurance training, whereas HSP27 is influenced only by endurance training, suggesting the existence of a training-modality-specific action on the adaptive processes including HSPs in human skeletal muscle (Folkesson et al. 2013). The function of HSP70 depends on ATP hydrolysis whereas that of HSP27 does not. Moreover, HSP70 is an early responsive protein whereas HSP27 is a late responsive one. Consequently, these proteins have been referred to as complementary protective proteins (Parcellier et al. 2003).

Basing on the gathered data on inflammatory response, the study was designed to evaluate the effect of 7-day training on HSP27 and HSP70 levels, and their relation to skeletal muscle damage and inflammation in elite canoeists.

Methods

Data collection and subjects. Six marathon canoeists, members of national team (Table 1), were observed during preparatory training period (pre-season, March) at the first, the fourth and after 7 days of the conditioning camp, and after 3 days of the recovery. The observations were performed at the Sport Centre in Gryfino (Poland). Details of the training program are presented in Table 2. All subjects occupied the same accommodations and followed the same training and diet schedules.

Table 1. Anthropometric and body composition data in canoeists; (mean ± SD)

	1 st day of camp n = 6	4 th day of camp n = 6	After 7-day of camp n = 6	After 3-day recovery n = 6
Height cm	181.2 ±9.8	181.2 ±9.8	181.2 ±9.8	181.2 ±9.8
BM kg	79.7 ±6.6	80.1 ±6.1	80.1 ±5.9	80.0 ±6.2
BMI kg/m ²	24.3 ±2.1	24.5 ±2.2	24.5 ±2.1	24.4 ±2.1
FFM kg	68.9 ±6.0	70.8 ±5.0	70.1 ±5.2	69.9 ±5.4
FM kg	10.8 ±2.7	9.3 ±3.2	10.0 ±3.1	10.0 ±3.0
FM%	13.5 ±3.0	11.6 ±3.5	12.4 ±3.3	12.5 ±3.2

BM – Body Mass; BMI – Body Mass Index; FFM – Fat-Free Mass; FM – Fat Mass.

Table 2. The 7-day training program in canoe during conditioning camp (March)

	a.m. 10:00–12:00	TRAINING INTENSITY	p.m. 16:00–18:00	TRAINING INTENSITY
1 st day of camp blood and body composition 7.00 and 8.00 a.m.	long distance paddling of 14 km (focus on technique)	moderate	repeat sets of 4 × 10/3' break; distance of 14 km long distance run of 10 km	moderate – high moderate
2 nd day of camp	repeat sets of 3 × (5 × 5/1'p)/5' break; distance of 20 km	moderate – high	repeat sets of 10 × 30"; distance of 12 km strength training 10 exercise; 6x10 reps (25 ton)	high moderate
3 rd day of camp	repeat sets of 3 × 30/5' break; distance of 20 km	moderate	rest; sauna	–
4 th day of camp blood and body composition 7.00 and 8.00 a.m.	repeat sets of 6 × 10/2' break; distance of 16 km	high	long distance paddling of 12 km (focus on technique) strength training; 10 exercise; 4 × 25 reps (30 ton)	moderate moderate
5 th day of camp	repeat sets of 3 × (2/1' + 4/2' + 8/3' + 4/2' + 2')/5' break; distance of 16 km	high	long distance paddling of 12 km (focus on technique) long distance run of 10 km	moderate moderate
6 th day of camp	repeat sets of 4 × 15/4' break; distance of 16 km	moderate	repeat sets of 10 × 30"; distance of 12 km strength training; 10 exercise; 6 × 10 reps (25 ton)	high moderate
7 th day of camp	long distance paddling of 12 km (focus on technique) long distance run of 8 km	moderate moderate	rest; sauna	–
END OF CAMP				
blood and body composition 7.00 and 8.00 a.m.				
AFTER 3-DAY RECOVERY				
blood and body composition 7.00 and 8.00 a.m.				

All the subjects were informed of the aim of the study and were given their written consent for participation in the project. In two cases, the consent was obtained from parents. The protocol of the study was approved by the ethics committee at Medical University Poznan, in accordance with the Helsinki Declaration.

Body composition. Body mass (BM) and body composition (fat-free mass FFM and fat mass FM) were estimated using a bioelectrical impedance (BIA) by using Tanita Body Composition Analyser MC-980 (Japan) calibrated prior to each test session in accordance to the manufacturer's guidelines. Duplicate measures were taken with the participant in a standing position; the average value was used for the final analysis. The recurrence of measurement was 98%. The measurements were taken between 7.00 and 8.00 a.m. before blood sampling.

Blood sampling. Blood samples were taken from the elbow vein between 7.00 and 8.00 a.m. after 15 minutes of body composition analysis. Within 20 min, blood samples were centrifuged at 1,000–2,000 x g and +4°C for 10 min. Aliquots of serum were stored at –80°C.

Skeletal muscle damage. Serum total creatine kinase (CK) activity was used as a marker of sarcolemma disruption and was evaluated by using commercially available reagents and Dr Lange analyser (Germany) at a temperature of 20–25°C. CK detection limit for the applied kit was 6 IU/L. The intra-assay coefficient of variation for the CK kit was <2%.

Inflammatory mediators. Serum interleukin-1 β (IL-1 β) and tumour necrosis factor α (TNF α) levels were evaluated by enzyme immunoassay methods using R&D Systems kits (USA). Detection limits for IL-1 β and TNF α were 0.023 pg/mL and 0.038 pg/mL, respectively. The average intra-assay coefficient of variation was about 8.0%.

Heat shock proteins. Serum heat shock proteins HSP27 and HSP70 were evaluated by enzyme immunoassay methods using Calbiochem kit (USA) and Stressgen kit (USA). Detection limits for HSP27 and HSP70 were 0.02 ng/mL and 0.2 ng/mL, respectively. Intra-assay coefficients of variation (CV) were <5% for both HSPs.

Statistical analysis. For all tested parameters the arithmetic mean (\bar{x}) and standard deviation (SD) were calculated. The conformity of distribution was tested with the Shapiro-Wilk test. The level of critical significance was set at $p < 0.05$. As the conditions of normal distribution or uniformity of variance were not met, non-parametric Friedman ANOVA test was used. The significance of differences between the dates of tests was assessed using a post-hoc Friedman ANOVA test. Statistical analysis was carried out using STATISTICA 10.0.

Results

Body composition. There were not significant changes in body composition but a tendency to high values in FFM. All subjects had normal BMI ranged from 21.5 to 27.7 kg/m².

Skeletal muscle damage (Table 3). The CK activity, as a marker of muscle damage, did not change significantly during the conditioning camp where canoe endurance training dominated.

Inflammatory mediators (Table 3). The concentrations of IL-1 β and TNF α did not change at the same time. IL-1 β level were significantly higher at the 1st day of camp whereas TNF α reached the highest level after 3 days of recovery. There was a negative correlation between IL-1 β and TNF α ($r = -0.490$, $P < 0.05$).

Table 3. The serum levels of creatine kinase (CK), cytokines IL-1 β and TNF α as well as heat shock proteins HSP27 and HSP 70 in canoeists; (mean \pm SD)

	1 st day of camp n = 6	4 th day of camp n = 6	After 7-day of camp n = 6	After 3-day recovery n = 6	Differences
CK IU/L	80.0 \pm 9.4	74.9 \pm 19.3	71.1 \pm 19.9	73.0 \pm 19.0	Chi ² ANOVA = 3.8 P = 0.283
IL-1 β pg/mL	1.24 \pm 0.09	1.19 \pm 0.16	0.98 \pm 0.09	1.04 \pm 0.10	Chi ² ANOVA = 12.6 P < 0.01
TNF α pg/mL	3.14 \pm 0.13	3.36 \pm 0.12	3.68 \pm 0.11	4.13 \pm 0.17	Chi ² ANOVA = 18 P < 0.001
HSP27 pg/mL	766 \pm 54	706 \pm 58	688 \pm 60	650 \pm 68	Chi ² ANOVA = 9.8 P < 0.05
HSP70 ng/mL	1.24 \pm 0.07	1.10 \pm 0.09	1.28 \pm 0.09	1.42 \pm 0.06	Chi ² ANOVA = 13.4 P < 0.01

Heat shock protein (Table 3). Similarly to cytokines, HSP27 and HSP70 concentrations did not change at the same time. HSP27 level was significantly higher at the 1st day of camp whereas HSP70 reached the highest level after 3 days of recovery. Figure 1 show the significant negative relation between TNF α and HSP27 ($r = -0.562$, $P < 0.01$), and positive between TNF α and HSP70 ($r = 0.651$, $P < 0.01$). It shows that the release of HSP27 and HSP70 into the circulation may occur in response to exercise-induced inflammation.




Figure 1. The significant relations between TNF α and HSP27 (top; $r = -0.563$, $P < 0.01$) as well as between TNF α and HSP70 (bottom; $r = 0.651$, $P < 0.01$)

Discussion

In response to exercise-induced muscle damage (EIMD), myogenic satellite cells become activated, proliferate, and repopulate the myofiber population by fusing together or fusing with existing myofibers. This process is mediated largely through various molecules such as cytokines and HSPs that participate in muscle regeneration and hypertrophy (Shortreed et al. 2008).

Disruption of the myofiber integrity is reflected by increased serum or plasma CK activity. In human and animal models, increased CK activity is observed after mechanical stress, e.g. extensive physical exercise, and in the course of muscle degenerative diseases (Brancaccio et al. 2010). Athletes, as a rule, have higher serum CK activity than non-athletes because of the regular strain imposed by training on their muscles. In the present study, CK did not reach the high activity after 7-day conditioning camp where the canoe endurance training dominated. It is commonly known that EIMD plays a role in promoting skeletal muscle hypertrophy (Flann et al. 2011; Sandri 2008; Shortreed et al. 2008). However, some researchers have questioned this hypothesis, noting that hypertrophy can occur in the relative absence of muscle damage. Although an eccentric exercise has greater hypertrophic effects compared with other types of actions, a cause-effect relationship directly linking these gains to EIMD is yet to be established. Moreover, if such a relationship does in fact exist, it is not clear to what extent of damage it is optimal for inducing maximum muscle growth (Schoenfeld 2012).

Although EIMD did not occur in marathon canoeists, both cytokines and HSPs concentrations changed following sport training. The serum concentrations of IL-1 β and HSP27 were considerably higher at the first day while concentrations of TNF α and HSP70 were higher after 7-day conditioning camp. It is very interesting that TNF α and HSP70 reached the highest levels after 3-day recovery.

In our study, IL-1 β and HSP27 constantly decreased after 7-day conditioning camp and after 3-day recovery. Data have shown that both molecules are expressed only in response to high-force eccentric exercise and are related to damaged myofibers (Fielding et al. 1993; Koh 2002; Paulsen et al. 2009). However, it does not seem likely that IL-1 β and HSP27 have any direct involvement in muscle hypertrophy. However, HSP27 has been proposed to play a direct role in protecting skeletal muscle from contraction-induced damage via interactions with cytoskeletal elements and in regulation of the glutathione system (Koh 2002).

TNF α is considered to play a major role in the regulation of muscle mass. One of the possible actions of TNF α on skeletal muscle is generation of reactive oxygen species and modification of signalling pathways. On the other hand, TNF α can directly induce muscle catabolism by the inhibition of protein synthesis and myogenesis in myoblasts. Moreover, studies have indicated that TNF α is also responsible for triggering the death receptor mediated pathways of myonuclear apoptosis playing a significant role in muscle loss (Fernández-Celemin et al. 2002). The pro-catabolic action of TNF α was accompanied by increase in HSP70 (Figure 1). More recently, the presence of muscle-derived HSP70 in the circulation in response to aerobic and eccentric exercises has been demonstrated (Febbraio et al. 2002; Heck et al. 2012; Thompson et al. 2002). However, further studies have not confirmed that the muscle could be the major source of circulatory HSP70 precluded the 'muscle hypothesis', and suggested that other tissues should be responsible for the increase of HSP70 into the circulation (Heck et al. 2012). The serum and muscle HSP70 concentrations are dependent on both training intensity and volume (Liu et al. 2000). The elevated level of HSP70 was observed in rowers, soccer players, endurance runners and tennis players (Banfi et al. 2006; Fehrenbach et al. 2000; Liu et al. 2000; Ziemann et al. 2013).

Analysis of body composition in canoeists did not show significant changes in FM and FFM, and any relation with cytokines and HSPs. Our previous study demonstrated that FM and FFM in canoeists are similar to healthy non-athletes at the same age but significantly lower compared with judoists and wrestlers. In combat sports, the high FFM and CK activity were accompanied by high IL-1 β and TNF α levels. This shows that type of sport discipline and training load markedly effect on relation of body composition with inflammatory response (Zembron-Lacny et al. 2013).

Conclusions

These results show that 7-day canoeing training modulates pro-inflammatory response which is related to HSPs release into the circulation, and reveal that skeletal muscle damage is not necessary to induce training-induced inflammation.

Acknowledgments

This study was financially supported by a grant, NRSA 1003151, from the Ministry of Science and High Education, Poland.

References

- Banfi G., Malavazos A., Iorio E., Dolci A., Doneda L., Verna R., Corsi M.M. Plasma oxidative stress biomarkers, nitric oxide and heat shock protein 70 in trained elite soccer players. *Eur J Appl Physiol.* 2006; 96: 483–486.
- Borges G.F., Rama L., Pedreiro S., Alves F., Santos A., Massart A., Paiva A., Teixeira A.M. Differences in plasma cytokine levels between elite kayakers and nonathletes. *Biomed Res Int.* 2013: 370354; DOI 10.1155/2013/370354.
- Brancaccio P., Lippi G., Maffulli N. Biochemical markers of muscular damage. *Clin Chem Lab Med.* 2010; 48: 757–767.
- Febbraio M.A., Ott P., Nielsen H.B., Steensberg A., Keller C., Krstrup P., Secher N.H., Pedersen B.K. Exercise induces hepatosplanchnic release of heat shock protein 72 in humans. *J Physiol.* 2002; 544: 957–962.
- Fehrenbach E., Niess A.M., Schlotz E., Passek F., Dickhuth H.H., Northoff H. Transcriptional and translational regeneration of heat shock proteins in leukocytes of endurance runners. *J Appl Physiol.* 2000; 89: 704–710.
- Fernández-Celemin L., Pasko N., Blomart V., Thissen J.P. Inhibition of muscle insulin-like growth factor I expression by tumor necrosis factor-alpha. *Am J Physiol Endocrinol Metab.* 2002; 283: 1279–1290.
- Fielding R.A., Manfredi T.J., Ding W., Fiatarone M.A., Evans W.J., Cannon J.G. Acute phase response in exercise. III. Neutrophil and IL-1 beta accumulation in skeletal muscle. *Am J Physiol.* 1993; 265: R166–R172.
- Flann K.L., LaStayo P.C., McClain D.A., Lindstedt S.L. Muscle damage and muscle remodeling: no pain, no gain? *J Exp Biol.* 2011; 214: 674–679.
- Folkesson M., Mackey A.L., Langberg H., Oskarsson E., Piehl-Aulin K., Henriksson J., Kadi F. The expression of heat shock protein in human skeletal muscle: effects of muscle fibre phenotype and training background. *Acta Physiol (Oxf).* 2013; DOI 10.1111/apha.12124.
- Heck T.G., Scomazzon S.P., Ludwig M.S., de Bittencourt P.I.Jr. Role of heat shock proteins in skeletal muscle. *Intech Open Science* 2012; <http://cdn.intechopen.com/pdfs-wm/38432.pdf>.
- Koh T.J. Do small heat shock proteins protect skeletal muscle from injury? *Exerc Sport Sci Rev.* 2002; 30: 117–121.
- Liu Y., Lormes W., Baur C., Optiz-Gress A., Altenburg D., Lehmann M., Steinacker J.M. Human skeletal muscle HSP70 response to physical training depends on exercise intensity. *Int J Sports Med.* 2000; 21: 351–355.
- Mackey A.L., Kjaer M., Dandanell S., Mikkelsen K.H., Holm L., Døssing S., Kadi F., Koskinen S.O., Jensen C.H., Schrøder H.D., Langberg H. The influence of anti-inflammatory medication on exercise-induced myogenic precursor cell responses in humans. *J Appl Physiol.* 2007; 103: 425–431.
- Main L.C., Dawson B., Grove J.R., Landers G.J., Goodman C. Impact of training on changes in perceived stress and cytokine production. *Res Sports Med.* 2009; 17: 121–132.

- Marin D.P, Santos Rde C., Bolin A.P., Guerra B.A., Hatanaka E., Otton R. Cytokines and oxidative stress status following a handball game in elite male players. *Oxid Med Cell Longev*. 2011; 804873; DOI 10.1155/2011/804873.
- Madamanchi N.R., Li S., Patterson C., Runge M.S. Reactive oxygen species regulate heat-shock protein 70 via the JAK/STAT pathway. *Arterioscler Thromb Vasc Biol*. 2001; 21: 321–326.
- Noble E.G., Milne K.J., Melling C.W.J. Heat shock proteins and exercise: a primer. *Appl Physiol Nutr Metab*. 2008; 33: 1050–1065.
- Parcellier A., Schmitt E., Gurbuxani S., Seigneurin-Berny D., Pance A., Chantôme A., Plenchette S., Khochbin S., Solary E., Garrido C. HSP27 is a ubiquitin-binding protein involved in I-kappaB α proteasomal degradation. *Mol Cell Biol*. 2003; 23: 5790–5802.
- Paulsen G., Lauritzen F., Bayer M.L., Ugelstad I., Owe S.G., Hallén J., Bergersen L.H., Raastad T. Subcellular movement and expression of HSP27, B-crystallin, and HSP70 after two bouts of eccentric exercise in humans. *J Appl Physiol*. 2009; 107: 570–582.
- Pedersen B.K. Muscles and their myokines. *J Exp Biol*. 2011; 214: 337–346.
- Reinke S., Karhausen T., Doehner W., Taylor W., Hottenrott K., Duda G.N., Reinke P., Volk H.D., Anker S.D. The influence of recovery and training phases on body composition, peripheral vascular function and immune system of professional soccer players. *PLoS ONE*. 2009; 4: e4910: 1–7.
- Roubenoff R. Physical activity, inflammation and muscle loss. *Nutr Rev*. 2007; 65: S208–S212.
- Sandri M. Signaling in muscle atrophy and hypertrophy. *Physiology*. 2008; 23: 160–170.
- Schoenfeld B.J. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J Strength Cond Res*. 2012; 26: 1441–1453.
- Schutz Y., Kyle U.G., Pichard C. Fat-free mass index and fat mass index percentiles in Caucasians aged 18–98 y. *Int J Obes*. 2002; 26: 953–960.
- Shortreed K., Johnston A., Hawke T.J. Satellite cells and muscle repair. In: *Skeletal Muscle Damage and Repair*. Ed. P.M. Tiidu). Human Kinetics. Champaign, IL, USA 2008: 77–88.
- Steinacker J.M., Lormes W., Reissnecker S., Liu Y. New aspects of hormone and cytokine response to training. *Eur J Appl Physiol*. 2004; 4: 382–391.
- Tayek J.A. Effects of tumour necrosis factor alpha on skeletal muscle amino acid metabolism studied in-vivo. *J Am Coll of Nutr*. 1996; 15: 164–168.
- Thompson H.S., Clarkson P.M., Scordilis S.P. The repeated bout effect and heat shock proteins: intramuscular HSP27 and HSP70 expression following two bouts of eccentric exercise in humans. *Acta Physiol Scand*. 2002; 174: 47–56.
- Tracey K.J., Wei H., Manogue K.R., Fong Y., Hesse D.G., Nguyen H.T., Kuo G.C., Beutler B., Cotran R.S., Cerami A. Cachectin/tumor necrosis factor induces cachexia, anemia, and inflammation. *J Expe Med*. 1988; 167: 1211–1227.
- Weissman C. The metabolic response to stress: an overview and update. *Anesthesiology*. 1990; 73: 308–327.
- Zembroń-Łacny A., Slowinska-Lisowska M., Ziemba A. Integration of the thiol redox status with cytokine response to physical training in professional basketball players. *Physiol Res*. 2010; 59: 239–245.
- Zembroń-Łacny A., Ziemann E., Kasperska A., Zurek P., Rynkiewicz M., Rynkiewicz T., Laskowski R., Hübner-Woźniak E. Association between cytokine activity and body composition in highly trained athletes. *Medicina dello Sport*. 2013; 66: 199–209.
- Ziemann E., Zembroń-Łacny A., Kasperska A., Antosiewicz J., Grzywacz T., Garszka T., Laskowski R. Exercise training-induced changes in inflammatory mediators and heat shock proteins in young tennis players. *J Sports Sci Med*. 2013; 12: 282–289.

Cite this article as: Morawin B., Rynkiewicz M., Orysiak J., Rynkiewicz T., Zembroń-Łacny A. Exercise training-induced changes in inflammatory mediators and heat shock proteins in canoeists. *Centr Eur J Sport Sci Med*. 2014; 8 (4): 27–34.