

# World News of Natural Sciences

An International Scientific Journal

WNOFNS 26 (2019) 80-95

EISSN 2543-5426

---

---

## Earthworm species as pointer for herbaceous flora distribution and health of soil in Onigambari Forest Reserve, Oyo state Nigeria

**O. M. Onafurume**

Forestry Research Institute of Nigeria, Ibadan, Oyo State, Nigeria

Email address: [onodinho24@gmail.com](mailto:onodinho24@gmail.com)

### ABSTRACT

Amongst various and diverse organisms, earthworms (*Annelida: Oligochaeta*) are the most important components of soil biota when it comes to soil formation and maintenance of soil structure and fertility. The presence of earthworms modifies the soil quality of the environment due to their burrowing and casting. This affects the activities of other organisms. Thus, they are also termed “ecosystem engineers”. However, while we begin to understand the role of surface casts, it is still unclear to what extent plants utilize subsurface casts. This work is a study of the floral community structure as influenced by surface cast so as to determine the functional groups of earthworms in the natural forest zone of Onigambari Forest Reserve. This reserve is located at latitude 07°25'N and longitude 3°53'E within the low and semi-deciduous forest belt of Nigeria. It is divided into two: natural and plantation forests. Herein, the simple random sampling technique was used to select 10 geo-referenced plots (15m × 15m each). Three 1.0m × 1.0m square quadrats were randomly laid on each plot to give 30 sampling points surveyed for data collection. Data were collected on floral composition, relative importance values (RIV), earthworm composition and diversity, and physicochemical components of the soil. Twenty eight herbaceous plants were enumerated. *Andropogon gayanus* had the highest relative importance value (RIV) of (20.499), while *Acroceras zizanoides* and *Platostoma africanum* had the least RIV (1.1782) amongst other four. Seven earthworm species were enumerated, with *Eisenia fetida* having the highest RIV (28.571), while *Octoclasion cyaneam* and *Lumbricum rubellus* had the least. The species richness of earthworms was high (0.8061) and evenly distributed (0.8405), but with a low diversity ( $H' = 1.772$ ). There are two functional groups of earthworm in the natural forest zone, seven were surface dwelling, while only *Lumbricus terrestris* was sub-surface dwelling. The health of flora and soil of the natural forest zone of Onigambari forest reserve is low. Prevention of outside influence on the forest zone, seeding and deposition of organic waste matter to the soil are strategies that would enhance abundance and diversity of earthworms in the natural forest zone.

**Keywords:** Natural forest, Earthworms, Soil fertility, Relative importance value, Floristic composition

## 1. INTRODUCTION

Among various and diverse organisms found in the soil, earthworms (*Annelida: Oligochaeta*) are the most important components of soil biota when it comes to soil formation and maintenance of soil structure and fertility (Curry, 2004). Earthworms (EWs) play a key role in soil nutrient dynamics by altering the soil physical, chemical and biological properties which is generally in synchrony with plant demand (Lavelle et al., 1989). Therefore, integrating earthworms into agricultural/forest management in order to increase yield has attracted more focus (Lavelle et al., 1989; Senapati et al., 2002).

The crop production is usually higher in the soil with high number of earthworms than no or less earthworms (Edwards and Bater, 1992; Elmer, 2012). From evolutionary point of view the earthworms (EWs) are very old species. They survived over a million years due to their ability to adjust to different environmental conditions. They live mostly in damp soil enriched with organic substances. They take breaths through the skin, and they are very sensitive to changes in temperature, they are light on touch. During the winter they bury themselves in deeper layer to protect its body from low temperature, and during the summer and dryness to protect itself from dehydration (Brusca and Brusca, 2003).

Their muscle system is built with circular (segments) and longitudinal muscles and with their shrinkage and spread the earthworms are able to move. The body of EWs is covered with small fluffs, which is important in environmental adjustment and for search of the food in the soil. The waste product of earthworm's diet enrich the soil with nutritive substances, which stimulate the growth of plants. However, the earthworms are very important source of diet for numerous animals in the soil. They are hermaphrodites, which means that the each individual has both female and male reproductive systems. This attribute also contributes well to their environmental adjustment, because it aides them to reproduce very easily. Their eggs are hatched in soil protected with capsule, which arises from the secret *clitellum* (front part). The capsule protects the young worm until complete maturation (Pechenik, 2009).

Investigating the potential of the earthworms to integrate into agricultural/forestry management, knowledge on different physical, chemical and management factors that affect the distribution and abundance of earthworm population is important as that will help to identify the ecological appropriateness of the earthworms in order to supplement their existing population and quantify the impact of earthworms on agricultural/forest lands (Mele and Carter, 1999). Not only from agricultural perspective, earthworms are equally important from ecological point of view because they contain highest soil macro-faunal biomass (Edwards and Bohlen, 1996) and are also increasingly regarded as bio-indicators of soil quality (Pères et al., 2011).

Earthworms (EWs) are the major decomposers of dead and decomposing organic matter, and acquire their nutrition from the bacteria and fungi that grow upon these materials. They decompose the organic matter and make the major contributions to recycling the containing nutrients. Earthworms occur in the warmest soils and many tropical soils. They are divided into 23 families, more than 700 genera, and more than 7,000 species. Their size ranges from an inch to two yards, and are found seasonally at all depths of the soil (Pechenik, 2009). The presence of earthworms modifies the soil quality of the environment due to their various activities like burrowing and casting which affect the activities of other organisms. So, they are also termed as "ecosystem engineers" (Kalu et al, 2015).

Earthworms (*Annelida: Oligochaeta*) deposit several tons per hectare of casts enriched in nutrients and/or arbuscular mycorrhizal fungi (AMF) and create a spatial and temporal soil heterogeneity that can play a role in structuring plant communities (Zaller *et al.*, 2007).

Earthworms incorporate organic materials into the soil hence enhancing plant root penetration and can increase the water infiltration rates up to 10-folds. Research in the temperate regions show that casts are heterogeneously distributed showing associations with certain plant species and thereby specifically stimulating their growth (Zaller *et al.*, 2007). Earthworms (EWs) impact plant communities both through the modification of soil chemical, physical and microbiological properties and through seed ingestion (Bityutskii *et al.*, 2002). Seed ingestion by earthworms might impact seed germination and seedling growth through two possible different mechanisms: the provision of a nutrient-rich cast substrate that benefits seed germination, i.e., the earthworm cast, and the alteration of the seed coat. The quality of this substrate is closely linked to the earthworm capacity to choose the soil and litter particles that they ingest, which tend to increase cast content in organic matter, and to its capacity to modify soil properties, mainly through an increase in mineralization. Cast properties depend on both the ingested soil type and the earthworm species and seedlings that emerge from casts likely respond to these soil properties. The impact of earthworms on plant growth differs with plant species and with soil properties (Eisenhauer and Scheu, 2008; Laossi *et al.*, 2009). However, while we have some understanding on interactions between earthworm surface casts and plant species (Decaens *et al.*, 2004), we know virtually nothing regarding the functional significance of the casts on pasture flora distribution and carbon stocking potentials of tropical forest zones. Agricultural soils can significantly contribute to the global greenhouse gas (GHG) exchange, but the contribution varies among the gases. For nitrous oxide (N<sub>2</sub>O), the emissions from agricultural soils account for 60% of the anthropogenic emissions (Smith and Read, 1997), whereas for methane (CH<sub>4</sub>, mineral agricultural soils are usually sinks as the aerobic topsoil favours methanotrophic bacteria (Hütsch, 2001). For carbon-dioxide (CO<sub>2</sub>, soils can be either sinks or sources, depending on the balance of carbon input and output. N<sub>2</sub>O emissions are mainly regulated by soil oxygen status, but also by the availability of nitrogen and organic carbon (Granli and Bøckman, 1994). The oxygen availability varies with soil structure and moisture and the potential for N<sub>2</sub>O emissions is greatest when the water-filled pore space (WFPS) is 60–70% (Davidson, 1991) as this enables both nitrification and DE nitrification. When the WFPS is above 70%, only DE nitrification takes place due to the shortage of oxygen and the dominating end product is the N<sub>2</sub> gas. Earthworm (EWs) casts and burrow linings have high microbial activity and more denitrifying bacteria than the bulk soil (Brown *et al.*, 2000) and the moist anaerobic environment in the earthworm gut can stimulate microbial N<sub>2</sub>O production (Karsten and Drake, 1997). Earthworms can increase micro aggregate formation and the stability of soil carbon (Fonte *et al.*, 2007; Six and Paustian, 2014).

In the majority of terrestrial ecosystems, earthworms are the most abundant animal biomass (Lavelle & Spain, 2001). Earthworms are typical ecosystem engineers as they have a large impact on soil structure, which is not necessarily associated with trophic relationships. For example, the tropical earthworm *Reginaldia omodeoi*, Sims, formerly known as *Millsonia anomala*, can ingest up to 30 times its own biomass of soil per day, but very little of the ingested organic matter is then assimilated (8%). Furthermore, little of the assimilated carbon is used in biomass production (6%); the remainder is respired (94%) during activity and physical modifications of the soil (Lamotte & Bourlière, 1978; Lavelle, 1978). In temperate ecosystems, earthworms also ingest large amounts of material (2–15% of organic matter inputs) (Whalen &

Parmelee, 2000) and expend much energy in their modification of the soil (74–91% of assimilated carbon is respired) (Petersen & Luxton, 1982). Reduced tillage and no-till increase the densities of anecic, deep-burrowing earthworms in arable fields (Whalen and Parmelee, 2000). In the temperate and boreal fields, this group is mainly represented by the dew-worm, *Lumbricus terrestris* L. (Chen, 2001). In Finland, *L. terrestris* is the second most common earthworm species in arable fields, lagging only behind *Aporrectodea caliginosa*. (Nieminen *et al.*, 2008), and has the typical positive association with non-inversion cultivation (Nuutinen *et al.*, 2011). It is a large earthworm, which efficiently forages on crop residues (Shuster *et al.*, 2002) and builds middens (i.e. small mounds of collected litter and surface castings) at the openings of its permanent burrows, often penetrating deeper than 1m (Nuutinen and Butt, 2003). Trees, herbs, floral, hedges diversities, and land use have an influence on the spatial distribution of earthworm species. However, this aspect is little accounted for in literatures due to the scarcity of studies on it.

### 1. 1. Objectives of the study

The geographical distribution of earthworms in Onigambari forest reserve is still very poorly known and much valuable information can be obtained by sampling the various earthworm species that can be found there. The broad goal of this study is to investigate the effect of earthworm cast on herbaceous flora distribution and Carbon Sequestration potential of top soil in the natural forest zone of Onigambari forest reserve.


The specific objectives of the study were to:

- 1) To determine floral community structure of the natural forest zone of Onigambari forest reserve as influenced by surface casts
- 2) To determine the functional group(s) of earthworms (EWs) at the natural forest zone of Onigambari forest reserve
- 3) To determine the carbon stocking potential of the natural forest zone of Onigambari forest reserve under the influence of earthworm casts.


## 2. METHOD AND INSTRUMENTS /EQUIPMENT USED

### 2. 1. Study area

Onigambari forest reserve is located on latitude 07°25'N and longitude 3°53'E within the low and semi-deciduous forest belt of Nigeria (Picture 1). The forest reserve stands on an elevation range of 185 m – 205 m above sea level (ASL). The reserve is divided into two: natural and plantation forests. The natural forest zone of the forest reserve (which will be used for this study) consist of indigenous species such as *Terminalia sp*, *Tripochiton scleroxylon*, *Irvingia garbonensis*, *Treculia africana*, *Chrysophyllum albidum*, *Artocarpus atilus*, *Xylopia aethiopica* and *Terapleyra tetraptera* among other tree species. It is also composed of species physiognomy replica of a tropical vegetation pattern that is peculiar to a lowland rainforest consisting of trees, shrubs and herbaceous flora while the plantation forest is made up of exotic species such as *Gmelina arborea* and *Tectona grandis*. Indigenous or settler farmers from different parts of the country live in several communities located around the forest reserve. The natural forest zone of the reserve has a perimeter of 2.55 km and an area of 14.476 hectares (Earth Point, 2019).



Picture 1. Picture showing the map of Onigambari forest reserve



Picture 2. Google earth showing the map of Onigambari forest reserve (within the yellow line is the natural forest zone)

The study area is characterized by climatic conditions peculiar of the tropical rainforest, with a mean annual rainfall of 1,200 mm, and monthly mean temperature of 26°C, relative humidity of 75% and sunshine of 3.77 hours (Table 1). The vegetation of the area is determined by a combination of maritime air mass blowing in from the Blight of Benin in the Atlantic Ocean to the south region of the country, and Continental air mass that blows in from the Sahara desert in the north. The general region of the study area experiences a bimodal rainfall type with its peaks in June and September.

**Table 1.** Agro-meteorological parameters for Onigambari forest reserve and environs in 2017

Parameter (n=50)	Value
Mean monthly temperature	27 °C
Mean annual rainfall	1,200 mm
Mean monthly relative humidity	75%
Mean monthly wind run	7.81 Km/day
Mean daily sunshine hours	3.77 hours

SOURCE: NIMET, 2017

## 2. 2. Floral sampling procedures

### Layout of sampling Area and Plots for Floristic sampling

Simple random sampling technique was used to select the plots used for the assessment of the herbaceous flora. Statistical sampling function of MS Excel 2013 was used for the random selection of 10 plots for the floristic assessment. Herbaceous flora was sampled using three 1.0 m x 1.0 m square quadrats randomly laid on each of 15 m X 15 m randomly using GPS- located grids. A total of 10 grids were selected for the survey to give 30 sampling points surveyed for data collection.

Google earth (2015) was used to complement the layout with an aerial view of the Onigambari forest reserve in conjunction with Garmin™ 12 *etrex* Vista H model Global positioning system was used to locate and mark the different points of the coverage area, while the compass was used to ensure accuracy in the direction of the points in the location.

## 2. 3. Species identification and enumeration

### Herbaceous species

The herbaceous flora occurring in each quadrat was identified following published herbaceous handbooks of (Akobundu and Agyakwa, 1998) and (Johnson, 1997). Where on-field identification was not possible, the specimens were collected and preserved for later identification.

## **2. 4. Data collection**

Data on floristic composition and land use types were collected as follows;

### **Herbaceous flora**

1. Species types by visual and use of standard flora
2. Count of individuals in the quadrats and plots for abundance, frequency, density and diversity indices calculations
3. Fresh litter biomass per plot

### **Collection of earthworm data**

Earthworms were extracted from the soil using 60g of mustard seed powder mixed with 350 ml of water, and the following data were collected;

1. Species composition per plot
2. Density of earth worm per plot
3. Dry weight of cast per plot
4. Nutrient composition of cast

### **Collection of Soil data**

1. Top soil sample was collected From (0 – 25) cm depth using a pre-calibrated hand trowel
2. Nutrient composition, including soil Carbon
3. Soil organic matter per plot

## **2. 5. Data analysis**

Raw data collected were arranged and analyzed for species abundance, diversity (Dominance, Species richness/Simpson index, Evenness index and Equitability index), Relative Importance Values (RIV), carbon sequestration; such as the health of the natural forest zone of Onigambari forest reserve was determined by assessment of species diversity.

Species diversity indices were computed using Paleontological statistics Software (PAST 2.14) (Hammer *et al.*, 2001). The following was taken into account from the diversity index analysis following (Hammer *et al* 2001):

### **Abundance**

Abundance measures the quantity or number of each species and families in an ecological system.

### **Dominance**

This is a measure of the prevalence of a particular species in relation to other species in an ecosystem. Dominance shows the species with superior competitive ability to others regarded as inferior based on competition. It is the most of population in a community and the effect on any population is mainly influenced by the species of high dominance (Bernstein, 1981; Drews, 1998). It usually ranges from zero to one in value where one signifies high dominance.

### **Simpson index**

It measures the number of individual species and their abundance in an ecosystem. It shows the richness of a species in a habitat in relation to total number of species present. Its value ranges from zero to one where zero means low and one means completely rich in species.

### **Evenness**

It is a spread rate of occurrence of a species in an ecosystem. It usually ranges from zero to one with one signifying complete evenness of occurrence of species in an ecosystem.

### **Equitability index**

This is a measure of occurrence of a species or otherwise of species in an ecosystem. It ranges from zero to one.

### **Shannon Weiner**

This is a measure proposed by Claude Shannon and accounts for both abundance and evenness of species present in an ecosystem (Kent and Coker, 1992 and Olubode *et al.*, 2011). It measures overall community characteristics. It usually ranges from one to infinity where two and above signifies high random of species occurrence in an ecosystem.

### **Relative Importance Values (RIV)**

The Relative Importance Value (RIV) of all species were determined following (Olubode *et al.*, 2011). It was computed as:

$$\text{RIV} = \text{Relative frequency/Relative Density} \times 100$$

where:

### **Frequency**

The number of occurrence of a specie in a set of quadrats or area

### **Relative Frequency**

Is a relative value of occurrence of a specie in a set of quadrat to total species in the quadrats:

$$\text{Relative frequency} = \frac{\text{Frequency of a specie}}{\text{frequency of other species}} \times 100$$

### **Density**

Is the quantity of individual specie to abundance of species per unit area

$$\text{Density} = \frac{\text{Quantitative Values of a specie}}{\text{Quadrat size} \times \text{Number of quadrats laid}}$$

### Relative Density

This is the relative number of individual of specie. Relative density was obtained using the formula;

$$\text{Relative Density} = \frac{n_i}{N} \times 100$$

where:

$n_i$  is the total number of individual specie

$N$  is the total number of different species in the entire population.

### 3. RESULTS

#### Floristic compositions, herbaceous diversities and relative importance value of herbaceous flora in the study area

A total of twenty eight (28) species of herbaceous flora belonging to eleven (11) families were enumerated in the assessment of (30) quadrats laid in ten (10) plots. *Andropogon gayanus* (family: Asteraceae) had the highest relative importance value (20.499) in the study area. This was followed by *Desmodium scorpiurus* (family: Fabaceae) (15.783). Among species with the intermediate relative importance values are *Iisca rygoe*, *Shrankia leptocarpa*, *Ipomoaea triloba* and their RIVs respectively were 2.741, 2.33 and 2.33 (Table 2).

**Table 2.** Composition and relative importance values of flora in the study area.

S/N	Species	Relative Importance value
1	<i>Andropogon gayanus</i>	20.49939
2	<i>Desmodium scorpiurus</i>	15.78266
3	<i>Tithonia diversifolia</i>	6.07326
4	<i>Oplismenus burmannii</i>	4.713065
5	<i>Synedrella nodiflora</i>	4.688645
6	<i>Sclerocarpus africanus</i>	4.304029
7	<i>Mitracarpus villosus</i>	4.304029
8	<i>Cleome ruidosperma</i>	3.510379
9	<i>Chromolaena odorata</i>	3.125763
10	<i>Iisca rygo</i>	2.741148
11	<i>Commelina erecta</i>	2.741148

12	<i>Shrankia leptocarpa</i>	2.332112
13	<i>Ipomoaea triloba</i>	2.332112
14	<i>Bidens pilosa</i>	2.332112
15	<i>Tridax procumbens</i>	1.971917
16	<i>Senna occidentalis</i>	1.947497
17	<i>Mellanthera scandens</i>	1.947497
18	<i>Croton lobata</i>	1.947497
19	<i>Centrosema pubesens</i>	1.947497
20	<i>Sphenoclea zeylanica</i>	1.562882
21	<i>Poulzozia guineensis</i>	1.562882
22	<i>Platstomao africanum</i>	1.562882
23	<i>Schwenckia Americana</i>	1.178266
24	<i>Solanum nigrum</i>	1.178266
25	<i>Sida acuta</i>	1.178266
26	<i>Sagitaria latifolia</i>	1.178266
27	<i>Calopogonum mucunoides</i>	1.178266
28	<i>Acrocera zizanoides</i>	1.178266

*Eisenia fetida*, *Lumbricus terrestris*, *Eisenia hortensis*, *Allogophora chlorotica*, *Apporectodea longa*, *Lumbricum rubellus*, *Octoclasion cyaneam* were enumerated at the study area. *Eisenia fetida* had the highest relative density (28.57) at the study area, closely followed by *Lumbricus terrestris* and *Apporectodea longa* with relative densities of 21.43. *Eisenia hortensis*, *Allogophora chlorotica*, *Octoclasion cyaneam*, *Lumbricum rubellus* had relative densities of 7.14 (Table 3). Diversity indices (Table 4) of the earthworm indicated 7 species with a low dominance value (0.1939) across the study area, but with high species richness (Simpson = 0.8061). The species were highly evenly distributed in the ecosystem (0.8405), but on the overall, the diversity of the earthworms was quite low ( $H' = 1.772$ ). The pH of topsoil was higher 7.2, while the worm cast pH was relatively low 7.1. The values of nutrients in the worm cast were such that Nitrogen (N) = 2.15 – 0.67; Available Phosphorus (Avail P) = 8.14 – 7.29; Organic Carbon (Org C) = 22.39 – 3.66; and Potassium (K) = 0.24 – 0.60. However, the following were found to record increase after soil casting; Calcium (Ca) = 2.82 – 6.96; Magnesium (Mg) = 1.77 – 4.45; Sodium (Na) = 0.56 – 1.12; Manganese (Mn) = 108.6 – 301; Iron (Fe) = 52.3 – 145.0; Copper (Cu) = 3.10 – 24.90; and Zinc (Zn) = 2.44 – 6.0 (Table 5).

It is observed that worm cast had higher clay content (126 mg/kg) than top soil (86 mg/kg) as well as in sand (840 mg/kg) compared to 770 mg/kg in top soil. The textural class of the worm cast was loamy sand, while it was sandy loam for the topsoil of the study area (Table 6).

**Table 3.** Species composition of earthworms in the study area.

SN	Scientific name	Common name	Relative Density
1	<i>Eisenia fetida</i>	Common red worm	28.57143
2	<i>Lumbricus terrestris</i>	Dew worm	21.42857
3	<i>Eisenia hortensis</i>	night crawler	7.142857
4	<i>Allogophora chlorotica</i>	Green worm	7.142857
5	<i>Apporectodea longa</i>	Blackhead worm	21.42857
6	<i>Octoclasion cyaneam</i>	Blue-grey worm	7.142857
7	<i>Lumbricum rubellus</i>	Red earthworm	7.142857

**Table 4.** Diversity of earthworms in the study area

Indices	Value
Taxa_S	7
Dominance_D	0.1939
Simpson_1-D	0.8061
Shannon_H	1.772
Evenness_e <sup>H/S</sup>	0.8405

**Table 5.** Chemical properties of the Soil.

SN	Soil Organic matter	Top soil	Cast
1	pH (H <sub>2</sub> O)	7.2	7.1
2	Org C (g1kg)	22.39	3.66
3	N (g1kg)	2.15	0.67
4	Avail P (mg1kg)	8.14	7.29
5	Ca	2.82	6.96
6	Mg (Cawl1kg)	1.77	4.45
7	K	0.24	0.60
8	Na	0.56	1.12

<b>9</b>	Mn	108.6	301.0
<b>10</b>	Fe	52.3	145.0
<b>11</b>	Cu	3.10	24.90
<b>12</b>	Zn	2.44	6.0

**Table 6.** Distribution of Soil particles.

SN	Soil particles	Top soil	Cast
<b>1</b>	Exch (Acid)	0.6	0.4
<b>2</b>	Exch (H <sup>+</sup> )	0.6	0.4
<b>3</b>	Exch (H <sup>++</sup> )	0.0	0.0
<b>4</b>	Clay	86.0	126.0
<b>5</b>	Silt	144.0	34.0
<b>6</b>	Sand	770.0	840.0
<b>7</b>	Textural Class	Sandy loam	Loamy Sand

#### 4. DISCUSSION AND CONCLUSION

This study investigated the effect of earthworm cast on herbaceous flora distribution and carbon sequestration potential of top soil in the natural forest zone of Onigambari forest reserve. The study indicated that the abundance and diversity of Earthworms was not high. It was further observed that the fertility of the topsoil is related to some grazing activities. Contrary to what was expected, the topsoil was not very rich in fertility. This may be due to inadequate return of organic matter to the soil by the grazing animals. Moderate grazing and trampling usually increase the diversity of plants by decreasing the ability of any one plant species to become dominant and exclude other species (Rikhari et al., 1993; Singh et al., 2003). Thus, the natural forest zone of the study area was not properly monitored as it is observed that grazing animals such as cattle visit the forest reserve to browse through for pastures but sparsely find the right one to satisfy them in adequate quantity or some certain flora species have been overgrazed by this animals thereby reducing the organic waste matter that could be deposited back to the soil for earthworm activities, this in return affects the fertility of the soil negatively. More so, the wandering of the animals on the land affects the activities of earthworm on the soil as they are inhibited from efficient burrowing and surface casting. This was further supported by the many different plant species in various families with low similar relative importance values as indicated by findings of the study. *Andropogon gayanus* was the dominant species in the study area. This study shed more light on the importance of earthworms as indicators of health of the forest zone. Earthworm communities nearly always include species that pursue different ecological strategies and a familiarity with these strategies is essential to an understanding of

the structure of earthworm's communities. Earthworm casting activity has been reported to cause spatially and temporally heterogeneous soil resources which can be specifically utilized by various plant species (Jackson, 1992), eventually affecting the structure of plant communities (Zallar *et al.*, 2007). The earthworm enumerated were mainly surface functional groups, implicated in soil mineralization and litter processing. (Tomati *et al.* 1990) have reported positive correlation between top soil dwelling earthworms with surface casts and grasses. Only *Lumbricus tenerstris* is known to be subsurface dwelling (Zaller *et al.*, 2007) of all earthworms encountered. The natural forest zone of Onigambari forest was reserved with the aim of encouraging and protecting nature's biodiversity ranging from various tree species, various floristic composition in terms of various families and species of herbs and shrubs. However the study has shown that the aim of reserving the natural forest is not fully achieved as it is indicated to have *Andropogon gayanus* and *Desmodium scorpiurus* as the dominant specie with a very wide margin in their relative importance value from other floristic composition of the forest zone. Any management practice that would improve the abundance and diversity of the surface-dwelling functional groups of earthworms would help in recovery of the soil fertility in the natural forest zone of Onigambari forest reserve.

## **5. RECOMMENDATION**

This study has helped to identify a problem that probably may have not been taken into awareness by the natural forest conservationist of the study area which includes poor soil health and fertility, low earthworm abundance/diversity and poor flora distribution in terms of their relative importance value. Hence this has led this study to propose the following recommendations;

1. Proper monitoring of the forest by the forest guards to prevent outside influence on the natural forest zone of Onigambari forest reserve.
2. Strategies that will enhance abundance and diversity of earthworms in the forest zone e.g. seeding and deposition of organic waste matter to the soil.

## **References**

- [1] Akobundu, I. O. & Agyakwa C. W. (1998). *A Handbook of West African Weeds* (2nd edition.). International Institute of Tropical Agriculture, Ibadan, Oyo State, Nigeria.
- [2] Bernstein, B.B., Williams, B.E & Mann, K.H. 1981. The role of behavioural responses to predators in modifying urchins (*Strongylocentrotus droebachiensis*) destructive grazing and seasonal foraging patterns. *Mar. Biol.* 63: 39-49
- [3] Bityutskii, N.P., Lapshina, I.N., Lukina, E.I., Solov'eva, A.N., Patsevich, V.G. & Vygovskaya, A.A. 2002. Role of earthworms in mineralization of organic nitrogen compounds in soil. *Eurasian Soil Science*, 35, 1100–1107
- [4] Brown, G.G., Barois, I. and Lavelle, P., 2000. Regulation of soil organic matter dynamics and microbial activity in the drilosphere and the role of interactions with other edaphic functional domains. *European Journal of Soil Biology* 36: 177–198

- [5] Brusca RC, Brusca GJ (2003). Sinaur Associates, Sunderland, MA. *Journal of Invertebrates*, ISJ 10: 38-45, ISSN 1824-307X.
- [6] Chan, K.Y., Baker, G.H., Conyers, M.K., Scott, B. & Munro, K. 2004. Complementary ability of three European earthworms (Lumbricidae) to bury lime and increase pasture production in acidic soils of south-eastern Australia. *Applied Soil Ecology*, 26, 257–271
- [7] Chen H. Anti-tumor effect of earthworm extracts EE2. *Chin. Clin. Oncol.* 6: 349-350, 2001.
- [8] Curry JP (2004). Factors Affecting the Abundance of Earthworms in Soil. In CA Edwards (Ed). *Earthworm Ecology Second Edition*. CRC Press LLC. pp. 91-114.
- [9] Davidson, M.G. 1991. Equine ophthalmology. In, Gelatt, K.N. (Ed.), *Veterinary Ophthalmology*. 2nd ed. Lea & Febiger, Philadelphia, pp: 576-610.
- [10] Decaëns T., Jiménez J.J., Barros E., Chauvel A., Blanchart E., Fragoso C. & Lavelle P., 2004. Soil macrofaunal communities in permanent pastures derived from tropical forest or savanna. *Agr. Ecosyst. Environ.* 103: 301-312
- [11] Drews A.W. 1998. *Manual on Hydrocarbon analysis (6th edition)* ASTM International, West Conshohocken: Pennsylvania.
- [12] Edwards CA, Bater JE (1992). The use of earthworms in environmental management. *Soil Biol. Biochem.* 24: 1683-1689.
- [13] Edwards, C.A., Bohlen, P.J. (1996) *Biological and ecology of earth worms*. Chapman and Hall, London pp. 426
- [14] Eisenhauer, N. & Scheu, S. 2008. Earthworms as drivers of the competition between grasses and legumes. *Soil Biology and Biochemistry*, 40, 2650–2659
- [15] Elmer WH (2012). *Using Earthworms to Increase Soil Health and Suppress Diseases*. The Connecticut Agricultural Experiment Station.
- [16] Fonte, S.J. & Six, J. 2007. Earthworms and litter management contributions to ecosystem services in a tropical agroforestry system. *Ecological Applications*, 20, 1061–1073
- [17] Granli T. and Bockman O. C. (1994). Nitrous oxide from agriculture. *Norw. J. agric. Sci. (Suppl.)* 12: 1–128
- [18] Hammer, O., Harper, D. A. T and Ryan, P. D. 2001. Paleontological statistics software package for education and data analysis. *Paleontological Electronica* 4 (1) (4): 9
- [19] Hutsch, B (2001). Methane Oxidation in non-flooded Soils as affected by Crop Production – invited paper. *European Journal of Agronomy* 14, 237–260
- [20] Jackson C.R. (1992) *Characterization of subsurface flow on shallow-soiled hillslopes*. PhD dissertation, University of Washington, Seattle, WA.
- [21] Johnson I. (1997): *The useful plants of West Africa*, vol. 4 The Whitefriars Press Limited, Tonbridge, Kent TN9 1QR, Great Britain.

- [22] Karsten, G.R. & Drake, H.L. 1997. Denitrifying bacteria in the earthworm gastrointestinal tract and *in vivo* emission of nitrous oxide (N<sub>2</sub>O) by earthworms. *Applied and Environmental Microbiology*, 63, 1878–1882
- [23] Kent M and Coker P 1992 Vegetation Description and Analysis. A practical Approach. John Willey and Sons, New York, pp. 319.
- [24] Lamotte, M. & Bourlière, F. (eds) 1978. Problèmes d'écologie: structure et fonctionnement des écosystèmes terrestres. Masson, Paris.
- [25] Laossi, K.-R., Ginot, A., Noguera, D., Blouin, M. & Barot, S. 2009 a. Earthworm effects on plant growth do not necessarily decrease with soil fertility. *Plant and Soil* 328, 109–118
- [26] Laossi, K.-R., Noguera, D.C., Bartolomé-Lasa, A., Mathieu, J., Blouin, M. & Barot, S. 2009 b. Effects of endogeic and anecic earthworms on the competition between four annual plants and their relative reproduction potential. *Soil Biology and Biochemistry*, 41, 1668–1673
- [27] Lavelle, P. 1978. Les vers de terre de la savane de Lamto (Côte d'Ivoire): peuplements, populations et fonctions dans l'écosystème. Thèse d'état, Université Paris 6, Paris.
- [28] Lavelle P, Barois I, Martin A, Zaidi Z, Schaefer R (1989). Management of earthworm populations in agro-ecosystems: A possible way to maintain soil quality? In M Clarholm, L Bergström (Eds). Ecology of Arable Land - Perspectives and Challenges. Springer Netherlands. pp. 109-122.
- [29] Lavelle, P., Bignell, D., Lepage, M., Wolters, V., Roger, P., Ineson, P., *et al* 1997. Soil function in a changing world: the role of invertebrate ecosystem engineers. *European Journal of Soil Biology*, 33, 159–193
- [30] Lavelle, P. & Spain, A.V. 2001. *Soil Ecology*. Kluwer Scientific Publications, Amsterdam.
- [31] Mele PM, Carter MR (1999). Species abundance of earthworms in arable and pasture soils in south-eastern Australia. *Appl. Soil Ecol.* 12: 129-137
- [32] Nieminen, K., Immanen, J., Laxell, M., Kauppinen, L., Tarkowski, P., Dolezal, K., Tähtiharju, S., Elo, A., Decourteix, M., Ljung, K., Bhalerao, R., Keinonen, K., Albert, V.A., and Helariutta, Y. (2008). Cytokinin signaling regulates cambial development in poplar. *Proceedings of the National Academy of Sciences* 105: 20032-20037
- [33] Nuutinen, V. & Butt, K.R. 2003. Interaction of *Lumbricus terrestris* L. burrows with field subdrains. *Pedobiologia*, 47, 578–581
- [34] Nuutinen, V., Butt, K.R. & Jauhiainen, L. 2011. Field margins and management affect settlement and spread of an introduced dew-worm (*Lumbricus terrestris* L.) population. *Pedobiologia*, 54, S167–S172
- [35] Olubode, O.S., Awodoyin, R.O. and Ogunyemi, S. 2011. Floral diversity in the wetlands of Apete River, Eleyele Lake and Oba dam in Ibadan, Nigeria: Its implication for biodiversity erosion. *West African Journal of Applied Ecology* 18: 109-119

- [36] Pechenik JA. (2009). *Biology of the Invertebrates*. 5th edition McGraw-Hill Higher Education, Boston.
- [37] Pérès G, Vandenbulcke F, Guernion M, Hedde M, Beguiristain T, Douay F, Houot S, Piron D, Richard A, Bispo A, Grand C, Galsomies L, Cluzeau D (2011). Earthworm indicators as tools for soil monitoring, characterization and risk assessment. An example from the national Bioindicator programme (France). *Pedobiologia* 54, Supplement: S77-S87.
- [38] Petersen, H. & Luxton, M. 1982. A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos*, 39, 287–388
- [39] Rikhari HC, Negi GCS, Ram J, Singh SP (1993). Human-induced secondary succession in alpine meadow of Central Himalaya, India. *Arctic Alpine Res.* 25: 8-14
- [40] Schuster E, Dunn-Coleman N, Frisvad JC, Van Dijck PW M (2002). On the safety of *Aspergillus niger* – a review. *Appl. Microbiol. Biotechnol.* 59: 426-435
- [41] Senapati BK, Lavelle P, Panigrahi PK, Giri S, Brown G (2002). Restoring soil fertility and enhancing productivity in Indian tea plantations with earthworms and organic fertilizers. In: Program, Abstracts and Related Documents of the International Technical Workshop on Biological Management of Soil Ecosystems for Sustainable Agriculture, Série Documentos. pp. 172-190
- [42] Singh YP, Kumar A, Rai JPN (2003). Species diversity as related to grazing pressure in alpine meadows of Nanda Devi Biosphere Reserve. Proc. Nat. Seminar on Biodiversity conservation and management, EPCO, Bhopal, pp. 147-153.
- [43] Six, J., and Paustian, K., 2014. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology & Biochemistry* 32: 2099–2103
- [44] Six, J., Elliott, E.T., Paustian, K., and Doran, J.W., 1998. Aggregation and soil organic matter accumulation in cultivated and native grassland soils. *Soil Science Society of America Journal* 62: 1367–1377
- [45] Smith, S.E., and Read, D.J., 1997. *Mycorrhizal Symbiosis*. Academic Press, San Diego, CA, USA, p. 605.
- [46] Subin Kalu, Madan Koirala and Udhab Raj Khadaka (2015). *Journal of Ecology and Natural Environment*, vol. 7(5), pp. 124-131. DOI: 10.5897/JENE2015.0511
- [47] Tomati U, Galli E, Grappelli A, Dihena G (1990). Effect of earthworm casts on protein synthesis in radish (*Raphanus sativum*) and lettuce (*Lactuca sativa*) seedlings. *Biol. fert. Soil*, 9: 288-289
- [48] Whalen, J.K. & Parmelee, R.W. 2000. Earthworm secondary production and N flux in agroecosystems: a comparison of two approaches. *Oecologia*, 124, 561–573.
- [49] Zaller JG (2007). Vermicompost as a substitute for peat in potting media: effects on germination, biomass allocation, yields and fruit quality of three tomato varieties. *Sci. Hort.* 112: 191-199